Absorption and Drug Development

Solubility, Permeability, and Charge State

Alex Avdeef

CONTENTS

PREFACE xiii

ACKNOWLEDGMENTS xvii

DEFINITIONS ixx

1 INTRODUCTION 1

1.1 Shotgun Searching for Drugs? / 1
1.2 Screen for the Target or ADME First? / 2
1.3 ADME and Multimechanism Screens / 3
1.4 ADME and Medicinal Chemists / 4
1.5 The "A" in ADME / 5
1.6 It is Not Just a Number—It is a Multimechanism / 6

2 TRANSPORT MODEL 7

2.1 Permeability-Solubility-Charge State and the pH Partition Hypothesis / 7
2.2 Properties of the Gastrointestinal Tract (GIT) / 11
2.3 pH Microclimate / 17
2.4 Intracellular pH Environment / 18
2.5 Tight-Junction Complex / 18
2.6 Structure of Octanol / 19
2.7 Biopharmaceutics Classification System / 20

3 CHARGE STATE 22

3.1 Constant Ionic Medium Reference State / 23
3.2 pK_a Databases / 24
3.3 Potentiometric Measurements / 25
3.3.1 Bjerrum Plots / 25
3.3.2 pH Definitions and Electrode Standardization / 27
3.3.3 The "Solubility Problem" and Cosolvent Methods / 29
3.3.4 Use of Cosolvents for Water-Soluble Molecules / 30
3.4 Spectrophotometric Measurements / 31
3.5 Capillary Electrophoresis Measurements / 32
3.6 Chromatographic pK_a Measurement / 33
3.7 pK_a Microconstants / 33
3.8 pK_a "Gold Standard" for Drug Molecules / 35

4 PARTITIONING INTO OCTANOL 42

4.1 Tetrad of Equilibria / 43
4.2 Conditional Constants / 45
4.3 log P Databases / 45
4.4 log D / 45
4.5 Partitioning of Quaternary Ammonium Drugs / 50
4.6 log D of Multiprotic Drugs and the Common-Ion Effect / 50
4.7 Summary of Charged-Species Partitioning in Octanol–Water / 53
4.8 Ion Pair Absorption of Ionized Drugs—Fact or Fiction? / 53
4.9 Micro-log P / 54
4.10 HPLC Methods / 54
4.11 IAM Chromatography / 54
4.12 Liposome Chromatography / 55
4.13 Other Chromatographic Methods / 55
4.14 pH-Metric log P Method / 55
4.15 High-Throughput log P Methods / 59
4.16 Octanol–Water log P^N, log P^I, and log $D_{7.4}$ "Gold Standard" for Drug Molecules / 59

5 PARTITIONING INTO LIPOSOMES 67

5.1 Tetrad of Equilibria and Surface Ion Pairing (SIP) / 67
5.2 Databases / 69
5.3 Location of Drugs Partitioned into Bilayers / 69

5.4 Thermodynamics of Partitioning: Entropy- or Enthalpy-Driven? / 70
5.5 Electrostatic and Hydrogen Bonding in a Low-Dielectric Medium / 71
5.6 Water Wires, H^+/OH^- Currents, and the Permeability of Amino Acids and Peptides / 73
5.7 Preparation Methods: MLV, SUV, FAT, LUV, ET / 74
5.8 Experimental Methods / 75
5.9 Prediction of log P_{mem} from log P / 76
5.10 log D_{mem}, $diff_{mem}$, and the Prediction of log P^{SIP}_{mem} from log P^I / 79
5.11 Three Indices of Lipophilicity: Liposomes, IAM, and Octanol / 83
5.12 Getting it Wrong from One-Point log D_{mem} Measurement / 84
5.13 Partitioning into Charged Liposomes / 85
5.14 pK_a^{mem} Shifts in Charged Liposomes and Micelles / 86
5.15 Prediction of Absorption from Liposome Partition Studies? / 90
5.16 log P^N_{mem}, log P^{SIP}_{mem} "Gold Standard" for Drug Molecules / 90

6 SOLUBILITY **91**

6.1 Solubility–pH Profiles / 92
6.1.1 Monoprotic Weak Acid, HA (or Base, B) / 92
6.1.2 Diprotic Ampholyte, XH_2^+ / 93
6.1.3 Gibbs pK_a / 93
6.2 Complications May Thwart Reliable Measurement of Aqueous Solubility / 99
6.3 Databases and the "Ionizable Molecule Problem" / 100
6.4 Experimental Methods / 100
6.4.1 Saturation Shake-Flask Methods / 101
6.4.2 Turbidimetric Ranking Assays / 101
6.4.3 HPLC-Based Assays / 101
6.4.4 Potentiometric Methods / 101
6.4.5 Fast UV Plate Spectrophotometer Method / 107
6.4.5.1 Aqueous Dilution Method / 107
6.4.5.2 Cosolvent Method / 108
6.5 Correction for the DMSO Effect by the Δ-Shift Method / 111
6.5.1 DMSO Binding to the Uncharged Form of a Compound / 111
6.5.2 Uncharged Forms of Compound–Compound Aggregation / 112
6.5.3 Compound–Compound Aggregation of Charged Weak Bases / 112
6.5.4 Ionizable Compound Binding by Nonionizable Excipients / 113
6.5.5 Results of Aqueous Solubility Determined from Δ Shifts / 113

6.6 Limits of Detection / 115
6.7 log S_0 "Gold Standard" for Drug Molecules / 115

7 PERMEABILITY **116**

7.1 Permeability in the Gastrointestinal Tract and at the Blood–Brain Barrier / 116
7.2 Historical Developments in Artificial-Membrane Permeability Measurement / 118
7.2.1 Lipid Bilayer Concept / 118
7.2.2 Black Lipid Membranes (BLMs) / 123
7.2.3 Microfilters as Supports / 124
7.2.4 Octanol-Impregnated Filters with Controlled Water Pores / 128
7.3 Parallel Artificial-Membrane Permeability Assay (PAMPA) / 128
7.3.1 Egg Lecithin PAMPA Model (Roche Model) / 128
7.3.2 Hexadecane PAMPA Model (Novartis Model) / 129
7.3.3 Brush-Border Lipid Membrane (BBLM) PAMPA Model (Chugai Model) / 130
7.3.4 Hydrophilic Filter Membrane PAMPA Model (Aventis Model) / 131
7.3.5 Permeability–Retention–Gradient–Sink PAMPA Models (pION Models) / 131
7.3.6 Structure of Phospholipid Membranes / 131
7.4 The Case for the Ideal In Vitro Artificial Membrane Permeability Model / 132
7.4.1 Lipid Compositions in Biological Membranes / 132
7.4.2 Permeability–pH Considerations / 132
7.4.3 Role of Serum Proteins / 135
7.4.4 Effects of Cosolvents, Bile Acids, and Other Surfactants / 135
7.4.5 Ideal Model Summary / 137
7.5 Derivation of Membrane-Retention Permeability Equations (One-Point Measurements, Physical Sinks, Ionization Sinks, Binding Sinks, Double Sinks) / 137
7.5.1 Thin-Membrane Model (without Retention) / 139
7.5.2 Iso-pH Equations with Membrane Retention / 142
7.5.2.1 Without Precipitate in Donor Wells and without Sink Condition in Acceptor Wells / 143
7.5.2.2 Sink Condition in Acceptor Wells / 147

7.5.2.3 Precipitated Sample in the Donor Compartment / 147
7.5.3 Gradient pH Equations with Membrane Retention: Single and Double Sinks / 148
7.5.3.1 Single Sink: Eq. (7.34) in the Absence of Serum Protein or Sink in Acceptor Wells / 150
7.5.3.2 Double Sink: Eq. (7.34) in the Presence of Serum Protein or Sink in Acceptor Wells / 151
7.5.3.3 Simulation Examples / 152
7.5.3.4 Gradient pH Summary / 153
7.6 Permeability–Lipophilicity Relations / 153
7.6.1 Nonlinearity / 153
7.7 PAMPA: 50+ Model Lipid Systems Demonstrated with 32 Structurally Unrelated Drug Molecules / 156
7.7.1 Neutral Lipid Models at pH 7.4 / 160
7.7.1.1 DOPC / 166
7.7.1.2 Olive Oil / 167
7.7.1.3 Octanol / 168
7.7.1.4 Dodecane / 168
7.7.2 Membrane Retention (under Iso-pH and in the Absence of Sink Condition) / 169
7.7.3 Two-Component Anionic Lipid Models with Sink Condition in the Acceptor Compartment / 171
7.7.3.1 DOPC under Sink Conditions / 177
7.7.3.2 DOPC with Dodecylcarboxylic Acid under Sink Conditions / 179
7.7.3.3 DOPC with Phosphatidic Acid under Sink Conditions / 179
7.7.3.4 DOPC with Phosphatidylglycerol under Sink Conditions / 181
7.7.3.5 DOPC with Negative Lipids without Sink / 181
7.7.4 Five-Component Anionic Lipid Model (Chugai Model) / 181
7.7.5 Lipid Models Based on Lecithin Extracts from Egg and Soy / 183
7.7.5.1 Egg Lecithin from Different Sources / 183
7.7.5.2 Soy Lecithin and the Effects of Phospholipid Concentrations / 187
7.7.5.3 Lipophilicity and Decrease in Permeability with Increased Phospholipid Content in Dodecane / 194
7.7.5.4 Sink Condition to Offset the Attenuation of Permeability / 196

7.7.5.5 Comparing Egg and Soy Lecithin Models / 198
7.7.5.6 Titrating a Suspension of Soy Lecithin / 198
7.7.6 Intrinsic Permeability, Permeability–pH Profiles, Unstirred Water Layers (UWL), and the pH Partition Hypothesis / 199
7.7.6.1 Unstirred Water Layer Effect (Transport across Barriers in Series and in Parallel) / 199
7.7.6.2 Determination of UWL Permeability using pH Dependence (pK_a^{flux}) Method / 200
7.7.6.3 Determination of UWL Permeabilities using Stirring Speed Dependence / 205
7.7.6.4 Determination of UWL Permeabilities from Transport across Lipid-Free Microfilters / 207
7.7.6.5 Estimation of UWL Thickness from pH Measurements Near the Membrane Surface / 207
7.7.6.6 Prediction of Aqueous Diffusivities D_{aq} / 207
7.7.6.7 Intrinsic Permeability–log K_p Octanol–Water Relationship / 208
7.7.6.8 Iso-pH Permeability Measurements using Soy Lecithin–Dodecane–Impregnated Filters / 209
7.7.6.9 Gradient pH Effects / 211
7.7.6.10 Collander Relationship between 2% DOPC and 20% Soy Intrinsic Permeabilities / 215
7.7.7 Evidence of Transport of Charged Species / 215
7.7.7.1 The Case for Charged-Species Transport from Cellular and Liposomal Models / 218
7.7.7.2 PAMPA Evidence for the Transport of Charged Drugs / 221
7.7.8 Δ log P_e–Hydrogen Bonding and Ionic Equilibrium Effects / 222
7.7.9 Effects of Cosolvent in Donor Wells / 226
7.7.10 Effects of Bile Salts in Donor Wells / 228
7.7.11 Effects of Cyclodextrin in Acceptor Wells / 228
7.7.12 Effects of Buffer / 229
7.7.13 Effects of Stirring / 231
7.7.14 Errors in PAMPA: Intraplate and Interplate Reproducibility / 232
7.7.15 UV Spectral Data / 233
7.8 The Optimized PAMPA Model for the Gut / 236
7.8.1 Components of the Ideal GIT Model / 236

7.8.2 How Well Do Caco-2 Permeability Measurements Predict Human Jejunal Permeabilities? / 238

7.8.3 How Well Do PAMPA Measurements Predict the Human Jejunal Permeabilities? / 239

7.8.4 Caco-2 Models for Prediction of Human Intestinal Absorption (HIA) / 242

7.8.5 Novartis max-P_e PAMPA Model for Prediction of Human Intestinal Absorption (HIA) / 244

7.8.6 *p*ION Sum-P_e PAMPA Model for Prediction of Human Intestinal Absorption (HIA) / 244

8 SUMMARY AND SOME SIMPLE RULES **247**

REFERENCES **250**

INDEX **285**