



WILEY

CATALYSTS FOR FINE CHEMICAL SYNTHESIS

Regio- and Stereo-
Controlled Oxidations
and Reductions

5

Editors: Stanley M. Roberts and John Whittall

Contents

Series Preface	xvii
Preface to Volume 5	xix
Abbreviations	xxi
1 Industrial Catalysts for Regio- or Stereo-Selective Oxidations and Reductions A Review of Key Technologies and Targets	1
<i>John Whittall</i>	
1.1 Introduction	2
1.2 Reduction of Carbon-Carbon Double Bonds	3
1.2.1 Privileged structures: α -amino acids and itaconic acids	4
1.2.2 β -Amino acids	5
1.2.3 α -Alkyl substituted acids	6
1.2.4 α -Alkoxy substituted acids	8
1.2.5 Unsaturated nitriles	9
1.2.6 Alkenes and allyl alcohols	10
1.2.7 α,β -Unsaturated aldehyde reduction	10
1.3 Ketone and Imine Reduction	12
1.3.1 Catalytic hydrogenation of ketones and imines	12
1.3.2 Asymmetric transfer hydrogenation (ATH) catalysts	15
1.3.3 Modified borane reagents	20
1.3.4 Biocatalysts (alcohol dehydrogenases and ketoreductases)	21
1.4 Oxidation	23
1.4.1 Sharpless chiral epoxidation of allyl alcohols	23
1.4.2 Dioxirane catalyzed epoxidation	23
1.4.3 Amines and iminium salts	25
1.4.4 Phase transfer catalysts	25
1.4.5 The Juliá-Colonna method (polyleucine oxidation)	26
1.4.6 Organocatalytic α -hydroxylation of ketones	27
1.4.7 Baeyer–Villiger oxidation	27
1.4.8 Chiral sulfoxides	28
References	29
2 Asymmetric Hydrogenation of Alkenes, Enones, Ene-Esters and Ene-Acids	35
2.1 (<i>S</i>)-2,2'-Bis{[di(4-methoxyphenyl)phosphinyl]oxy}-5,5',6,6',7,7', 8,8'-octahydro-1,1'-binaphthyl as a ligand for rhodium-catalyzed asymmetric hydrogenation	36
<i>Ildiko Gergely, Csaba Hegedüs and Jozsef Bakos</i>	

2.1.1	Synthesis of (<i>S</i>)-5,5',6,6',7,7',8,8'-Octahydro-1,1'-bi-2-naphthol . . .	37
2.1.2	Synthesis of (<i>S</i>)-2,2'-Bis{[di(4-methoxyphenyl)phosphinyl]oxy}-5,5',6,6',7,7',8,8'-octahydro-1,1'-binaphthyl.	38
2.1.3	Asymmetric hydrogenation of Dimethyl itaconate.	40
	Conclusion.	41
	References.	41
2.2	Synthesis and application of phosphinite oxazoline iridium complexes for the asymmetric hydrogenation of alkenes.	42
	<i>Frederik Menges and Andreas Pfaltz</i>	
2.2.1	Synthesis of (4 <i>S</i> ,5 <i>S</i>)-2-(5-Methyl-2-phenyl-4,5-dihydro-oxazol-4-yl)-1,3-diphenyl-propan-2-ol	42
2.2.2	Synthesis of (4 <i>S</i> ,5 <i>S</i>)-O-[1-Benzyl-1-(5-methyl-2-phenyl-4,5-dihydro-oxazol-4-yl)-2-phenyl-ethyl]-diphenylphosphinite.	43
2.2.3	Synthesis of (4 <i>S</i> ,5 <i>S</i>)-[(η^4 -1,5-Cyclooctadiene)-{2-(2-phenyl-5-methyl-4,5-dihydro-oxazol-4-yl)-1,3-diphenyl-2-diphenylphosphinite-propane Iridium(I)}-tetrakis[3,5-bis(trifluoromethyl)phenyl]borate	45
2.2.4	Asymmetric hydrogenation of <i>trans</i> - α -Methylstilbene	46
	Conclusion.	47
	References.	48
2.3	Synthesis and application of heterocyclic phosphine oxazoline (HetPHOX) iridium complexes for the asymmetric hydrogenation of alkenes	48
	<i>Frederik Menges and Pier Giorgio Cozzi</i>	
2.3.1	Synthesis of (4 <i>S</i>)- <i>tert</i> -Butyl-2-(thiophene-2-yl)-4,5-dihydrooxazole	49
2.3.2	Synthesis of (4 <i>S</i>)- <i>tert</i> -Butyl-2-(3-diphenylphosphinothiophene-2-yl)-4,5-dihydrooxazole	50
2.3.3	Synthesis of (4 <i>S</i>)-[(η^4 -1,5-Cyclooctadiene)-{4- <i>tert</i> -butyl-2-(3-diphenylphosphinothiophene-2-yl)-4,5-dihydrooxazole} iridium(I)-tetrakis[3,5-bis(trifluoromethyl)phenyl]borate	52
2.3.4	Asymmetric hydrogenation of <i>trans</i> - α -Methylstilbene	53
	Conclusion.	54
	References.	54
2.4	(<i>R</i>)-2,2',6,6'-Tetramethoxy-bis[di(3,5-dimethylphenyl)phosphino]-3,3'-bipyridine [(<i>R</i>)-Xyl-P-Phos] as a ligand for rhodium-catalyzed asymmetric hydrogenation of α -dehydroamino acids.	55
	<i>Jing Wu and Albert S.C. Chan</i>	
2.4.1	Synthesis of 3-Bromo-2,6-dimethoxypyridine	55
2.4.2	Synthesis of Bis(3,5-dimethylphenyl)phosphine chloride	56
2.4.3	Synthesis of 3-Bromo-2,6-dimethoxy-4-di(3,5-dimethylphenyl)phosphinopyridine.	57
2.4.4	Synthesis of 3-Bromo-2,6-dimethoxy-4-di(3,5-dimethylphenyl)phosphinopyridine.	59
2.4.5	2,2',6,6'-Tetramethoxy-bis[di(3,5-dimethylphenyl)phosphinoyl]-3,3'-bipyridine	60
2.4.6	Optical resolution of (\pm)- 6 with (-) or (+)-2,3- <i>O,O'</i> -Dibenzoyltartaric acid monohydrate [(<i>R</i>)- 6 or (<i>S</i>)- 6]	61
2.4.7	(<i>R</i>)-2,2',6,6'-Tetramethoxy-bis[di(3,5-dimethylphenyl)phosphino]-3,3'-bipyridine [(<i>R</i>)-Xyl-P-Phos, (<i>R</i>)- 1]	62
2.4.8	Preparation of the stock solution of [Rh(<i>R</i> -Xyl-P-Phos)(COD)]BF ₄	63
2.4.9	A typical procedure for the asymmetric hydrogenation of methyl (<i>Z</i>)-2-Acetamidocinnamate.	64
	References.	65

2.5	(<i>R,R</i>)-2,3-Bis(<i>tert</i> -butylmethylphosphino)quinoxaline (QuinoXP*) as a ligand for rhodium-catalyzed asymmetric hydrogenation of prochiral amino acid and amine derivatives	65
	<i>Tsuneo Imamoto and Aya Koide</i>	
2.5.1	Synthesis of (<i>R</i>)- <i>tert</i> -Butyl(hydroxymethyl)methylphosphine–borane	66
2.5.2	Synthesis of (<i>R</i>)-Benzoyloxy(<i>tert</i> -butyl)methylphosphine–borane	67
2.5.3	Synthesis of (<i>S</i>)- <i>tert</i> -Butylmethylphosphine–borane	69
2.5.4	(<i>R,R</i>)-2,3-Bis(<i>tert</i> -butylmethylphosphino)quinoxaline (QuinoxP*)	70
2.5.5	Asymmetric hydrogenation of Methyl (<i>E</i>)-3-acetylamino-2-butenolate catalyzed by Rh(I)-(<i>R,R</i>)-2,3-Bis(<i>tert</i> -butylmethylphosphino)quinoxaline	71
	Conclusion	72
	References	73
2.6	Rhodium-catalyzed asymmetric hydrogenation of indoles	73
	<i>Ryoichi Kuwano and Masaya Sawamura</i>	
2.6.1	Synthesis of (<i>R</i>)-2-[(<i>S</i>)-1-(Dimethylamino)ethyl]-1-iodoferrocene	73
2.6.2	Synthesis of (<i>R</i>)-2-[(<i>S</i>)-1-(Diphenylphosphinyl)ethyl]-1-iodoferrocene	75
2.6.3	Synthesis of (<i>R,R</i>)-2,2'-Bis[(<i>S</i>)-1-(diphenylphosphinyl)ethyl]-1,1''-biferrocene	78
2.6.4	Synthesis of (<i>R,R</i>)-2,2''-Bis[(<i>S</i>)-1-(diphenylphosphino)ethyl]-1,1''-biferrocene [abbreviated to (<i>S,S</i>)-(<i>R,R</i>)-PhTRAP]	80
2.6.5	Catalytic asymmetric hydrogenation of <i>N</i> -Acetyl-2-butylindole	82
2.6.6	Catalytic asymmetric hydrogenation of 3-Methyl- <i>N</i> -(<i>p</i> -toluenesulfonyl)indole	84
	Conclusion	85
	References	86
3	Asymmetric Reduction of Ketones	87
3.1	(<i>R,R</i>)-Bis(diphenylphosphino)-1,3-diphenylpropane as a versatile ligand for enantioselective hydrogenations	89
	<i>Natalia Dubrovina and Armin Börner</i>	
3.1.1	Synthesis of (<i>S,S</i>)-1,3-Diphenylpropane-1,3-diol	89
3.1.2	Synthesis of (<i>S,S</i>)-Methanesulfonyloxy-1,3-diphenylpropane-1,3-diol	91
3.1.3	Synthesis of (<i>R,R</i>)-Bis(diphenylphosphino)-1,3-diphenylpropane	91
	Conclusion	93
	References	93
3.2	Synthesis of both enantiomers of 1-Phenylethanol by reduction of acetophenone with <i>Geotrichum candidum</i> IFO 5767	93
	<i>Kaoru Nakamura, Mikio Fujii and Yoshiteru Ida</i>	
3.2.1	Cultivation of <i>G. candidum</i> IFO 5767	94
3.2.2	Synthesis of (<i>S</i>)-1-Phenylethanol	95
3.2.3	Synthesis of (<i>R</i>)-1-Phenylethanol	95
	Conclusion	97
	References	97
3.3	Titanocene-catalyzed reduction of ketones in the presence of water. A convenient procedure for the synthesis of alcohols via free-radical chemistry	97
	<i>Antonio Rosales, Juan M. Cuerva and J. Enrique Oltra</i>	
3.3.1	Titanocene-catalyzed reduction of Acetophenone in the presence of water	98

3.3.2	Titanocene-catalyzed synthesis of Methyl 4-deuterio-4-phenyl-4-hydroxybutanoate	99
	References	100
3.4	Xyl-tetraPHEMP: a highly efficient biaryl ligand in the [diphosphine RuCl ₂ diamine]-catalyzed hydrogenation of simple aromatic ketones	101
	<i>Paul H. Moran, Julian P. Henschke, Antonio Zanotti-Gerosa and Ian C. Lennon</i>	
3.4.1	Synthesis of Tri(3,5-dimethylphenyl)phosphine oxide	102
3.4.2	Synthesis of Bis(3,5-dimethylphenyl)-(2-iodo-3,5-dimethylphenyl)phosphine oxide	103
3.4.3	Synthesis of <i>rac</i> -4,4',6,6'-Tetramethyl-2,2'-bis[bis(3,5-dimethylphenyl)phosphino]-biphenyl	105
3.4.4	Synthesis of <i>rac</i> -4,4',6,6'-Tetramethyl-2,2'-bis[bis(3,5-dimethylphenyl)phosphino]-biphenyl [abbreviated to (<i>rac</i>)-Xyl-tetraPHEMP]	106
3.4.5	Synthesis of [(<i>R</i>)- <i>N,N</i> -Dimethyl(1-methyl)benzylamino-C ² ,N]-{ <i>rac</i> -4,4',6,6'-tetramethyl-2,2'-bis[bis(3,5-dimethylphenyl)phosphino]-biphenyl}-palladium(II) tetrafluoroborate and separation of the diastereomers	107
3.4.6	Synthesis of (<i>S</i>)-4,4',6,6'-Tetramethyl-2,2'-bis[bis(3,5-dimethylphenyl)phosphino]-biphenyl: [abbreviated to (<i>S</i>)-Xyl-tetraPHEMP] and (<i>R</i>)-4,4',6,6'-Tetramethyl-2,2'-bis[bis(3,5-dimethylphenyl)phosphino]-biphenyl [abbreviated to (<i>R</i>)-Xyl-tetraPHEMP]	108
3.4.7	Synthesis of [(<i>R</i>)-Xyl-tetraPHEMP RuCl ₂ (<i>R,R</i>)-DPEN] and [(<i>S</i>)-Xyl-tetraPHEMP RuCl ₂ (<i>S,S</i>)-DPEN]	110
3.4.8	Reduction of Acetophenone using [(<i>S</i>)-Xyl-tetraPHEMP RuCl ₂ (<i>S,S</i>)-DPEN] as a precatalyst	111
	Conclusion	112
	References	112
3.5	<i>N</i> -Arenesulfonyl- and <i>N</i> -Alkylsulfamoyl-1,2-diphenylethylenediamine ligands for ruthenium-catalyzed asymmetric transfer hydrogenation of activated ketones	113
	<i>Michel (Massoud S.) Stephan and Barbara Mohar</i>	
3.5.1	Synthesis of <i>N</i> -Arenesulfonyl-1,2-diphenylethylenediamines	113
3.5.2	Preparation of Ru(II)- <i>N</i> -arenesulfonyl-1,2-diphenylethylenediamine complexes	114
3.5.3	Asymmetric transfer hydrogenation of Ethyl benzoylacetate	115
	Conclusion	116
	References	116
3.6	The synthesis and application of BrXuPHOS: a novel monodentate phosphorus ligand for the asymmetric hydrogenation of ketones	116
	<i>Martin Wills, Yingjian Xu, Gorden Docherty and Gary Woodward</i>	
3.6.1	Synthesis of (<i>S</i>)-BrXuPHOS	117
3.6.2	Synthesis of (<i>S,S,S,S</i>)-BrXuPHOS-Ru-DPEN	119
3.6.3	General procedure of asymmetric hydrogenation of acetophenone	120
	Conclusion	121
	Acknowledgement	121
	References	121
3.7	<i>In Situ</i> formation of ligand and catalyst: application in ruthenium-catalyzed enantioselective reduction of ketones	121
	<i>Jenny Wettergren and Hans Adolfsson</i>	
3.7.1	Synthesis of (<i>S</i>)-3-Fluoro-1-phenylethanol	122

	Conclusion	123
	References	124
3.8	Synphos and Difluorpos as ligands for ruthenium-catalyzed hydrogenation of alkenes and ketones	125
	<i>Séverine Jeulin, Virginie Ratovelomanana-Vidal and Jean-Pierre Genet</i>	
3.8.1	Synthesis of $[\text{RuCl}((S)\text{-SYNPHOS})(p\text{-cymene})]\text{Cl}$	125
3.8.2	Synthesis of $[\text{RuCl}((S)\text{-DIFLUORPHOS})(p\text{-cymene})]\text{Cl}$	126
3.8.3	Synthesis of $[\text{RuI}((S)\text{-DIFLUORPHOS})(p\text{-cymene})]\text{I}$	127
3.8.4	Synthesis of $[\text{NH}_2\text{R}_2] [(\text{RuCl}(\text{P}^*\text{P}))_2(\mu\text{-Cl})_3]$ $\text{P}^*\text{P} =$ SYNPHOS or DIFLUORPHOS and $\text{R} = \text{Me}$ or Et	127
3.8.5	Synthesis of $[\text{NH}_2\text{Me}_2][\text{RuCl}((S)\text{-DIFLUORPHOS})_2[\mu\text{-Cl}]_3]$	128
3.8.6	Synthesis of <i>in situ</i> generated $[\text{RuBr}_2((S)\text{-SYNPHOS})]$ and $[\text{RuBr}_2((S)\text{-DIFLUORPHOS})]$	129
	Conclusion	131
	References	131
3.9	An arene ruthenium complex with polymerizable side chains for the synthesis of immobilized catalysts	132
	<i>Estelle Burri, Silke B. Wendicke, and Kay Severin</i>	
3.9.1	Synthesis of 2-Methyl-cyclohexa-2,5-dienecarboxylic acid 2-(2-methyl-acryloyloxy)-ethyl ester	133
3.9.2	Synthesis of $[\eta^6\text{-}(2\text{-Methyl-benzoic acid } 2\text{-}(2\text{-methyl-acryloyloxy})\text{-ethyl ester})\text{RuCl}_2]_2$	134
	Conclusion	135
	References	135
3.10	Selective reduction of carbonyl group in β, γ -unsaturated α -alpha-ketoesters by transfer hydrogenation with Ru-(<i>p</i> -cymene) (TsDPEN)	135
	<i>Minjie Guo, Dao Li, Yanhui Sun and Zhaoguo Zhang</i>	
3.10.1	Synthesis of Di- μ -chloro-bis[chloro(η^6 -1-isopropyl-4-methyl-benzene)ruthenium(II)]	136
3.10.2	Synthesis of (\pm)-Monotosylate-1,2-diphenyl-1,2-ethylenediamine	136
3.10.3	Synthesis of Ru complex Ru(<i>p</i> -cymene)(TsDPEN)	138
3.10.4	Ru-TsDPEN catalyzed transfer hydrogenation reaction of β, γ -unsaturated- α -ketoesters	139
	Conclusion	140
	References	141
3.11	Preparation of polymer-supported Ru-TsDPEN catalysts and their use for the enantioselective synthesis of (<i>S</i>)-fluoxetine	141
	<i>Liting Chai, Yangzhou Li and Quanrui Wang</i>	
3.11.1	Synthesis of the supported ligand 9	141
3.11.2	Synthesis of ligand 17	148
3.11.3	General procedure for asymmetric transfer hydrogenation	150
3.11.4	Preparation of (<i>S</i>)-Fluoxetine hydrochloride	151
	Conclusion	154
	References	154
3.12	Polymer-supported chiral sulfonamide-catalyzed reduction of β -keto nitriles: a practical synthesis of (<i>R</i>)-Fluoxetine	155
	<i>Guang-yin Wang and Gang Zhao</i>	
3.12.1	Synthesis of (<i>R</i>)-3-Amino-1-phenyl-propan-1-ol	155
3.12.2	Synthesis of (<i>R</i>)-ethyl 3-hydroxy-3-phenylpropylcarbamate	156
3.12.3	Synthesis of (<i>R</i>)-3-(Methylamino)-1-phenylpropan-1-ol	157
3.12.4	Synthesis of (<i>R</i>)-Fluoxetine	158

Conclusion.....	159
References.....	159
4 Imine Reduction and Reductive Amination.....	161
4.1 Metal-free reduction of imines: enantioselective Brønsted acid-catalyzed transfer hydrogenation using chiral BINOL-phosphates as catalysts.....	162
<i>Magnus Rueping, Erli Sugiono, Cengiz Azap and Thomas Theissmann</i>	
4.1.1 Synthesis of (<i>R</i>)-2,2'-Bis-methoxymethoxy-[1,1'] binaphthalene (MOM-BINOL).....	162
4.1.2 Synthesis of (<i>R</i>)-3,3'-Diiodo-2,2'-bis(methoxymethoxy)-1,1'-binaphthalene.....	164
4.1.3 Synthesis of 3,3'-Bis-(3,5'-bis-trifluoromethyl-phenyl)-2,2'-bismethoxymethoxy [1,1'-binaphthalene].....	165
4.1.4 Synthesis of (<i>R</i>)-3,3'-[3,5-Bis(trifluoromethyl)phenyl]-1,1'-binaphthylphosphate.....	166
4.1.5 General procedure for the transfer hydrogenation of ketimines.....	167
4.1.6 Synthesis of [1-(2,4-Dimethyl-phenyl)-ethyl]- (4-methoxy-phenyl)-amine.....	167
Conclusion.....	168
References.....	170
4.2 Metal-free Brønsted acid-catalyzed transfer hydrogenation: enantioselective synthesis of tetrahydroquinolines.....	170
<i>Magnus Rueping, Thomas Theissmann and Andrey P. Antonchick</i>	
4.2.1 General procedure for the transfer hydrogenation of quinolines.....	170
4.2.2 Synthesis of 7-Chloro-4-phenyl-1,2,3,4-tetrahydroquinoline.....	172
4.2.3 Synthesis of (<i>S</i>)-2-Phenyl-1,2,3,4-tetrahydroquinoline.....	172
4.2.4 Synthesis of (<i>R</i>)-2-(2-(Benzo[1,3]dioxol-5-yl)ethyl)-1,2,3,4-tetrahydro-quinoline.....	173
Conclusion.....	174
References.....	174
4.3 A highly stereoselective synthesis of 3 α -Amino-23,24-bisnor-5 α -cholane via reductive amination.....	175
<i>Sharaf Nawaz Khan, Nam Ju Cho and Hong-Seok Kim</i>	
4.3.1 Synthesis of Tris[(2-ethylhexanoyl)oxy]borohydride.....	177
4.3.2 Synthesis of 3 α -Acetamino-23,24-bisnor-5 α -cholane.....	177
4.3.3 Synthesis of 3 α - <i>N</i> -1-[<i>N</i> (3-[4-Aminobutyl])-1,3-diaminopropane]-23,24-bisnor-5 α -cholane.....	179
Conclusion.....	181
Acknowledgements.....	181
References.....	181
5 Oxidation of Primary and Secondary Alcohols.....	183
5.1 Copper(II) catalyzed oxidation of primary alcohols to aldehydes with atmospheric oxygen.....	183
<i>Suribabu Jammi and Tharmalingan Punniyamurthy</i>	
5.1.1 Synthesis of copper(II) complex 1	184
5.1.2 Typical procedure for the oxidation of primary alcohols to aldehydes.....	185

	Conclusion	186
	References	187
5.2	Solvent-free dehydrogenation of secondary alcohols in the absence of hydrogen abstractors using Robinson's catalyst	187
	<i>G.B.W.L. Lighthart, R.H. Meijer, J. v. Buijtenen, J. Meuldijk, J.A.J.M. Vekemans and L.A. Hulshof</i>	
5.2.1	Dehydrogenation of 2-Octanol using Ru(OCOCF ₃) ₂ (CO)(PPh ₃) ₂ as a catalyst	187
	Conclusion	188
	References	188
5.3	2-Iodoxybenzoic acid (IBX)/ <i>n</i> -Bu ₄ NBr/CH ₂ Cl ₂ -H ₂ O: a mild system for the selective oxidation of secondary alcohols	188
	<i>Krisada Kittigowittana, Manat Pohmakotr, Vichai Reutrakul and Chutima Kuhakarn</i>	
5.3.1	Synthesis of 1-Hydroxy-5-decanone	189
	Conclusion	192
	References	192
6	Hydroxylation, Epoxidation and Related Reactions	193
6.1	Proline-catalyzed α -aminoxylation of aldehydes and ketones	194
	<i>Yujiro Hayashi and Mitsuru Shoji</i>	
6.1.1	Synthesis of (<i>R</i>)-2-Anilinoxypropanol	195
6.1.2	Synthesis of (<i>R</i>)-7-Anilinoxy-1,4-dioxaspiro[4.5]decan-8-one	196
	Conclusion	197
	References	198
6.2	Ru/Silia* Cat* TM TEMPO-mediated oxidation of alkenes to α -hydroxyacids	199
	<i>Rosaria Ciriminna and Mario Pagliaro</i>	
6.2.1	Synthesis of Silia*Cat* TM TEMPO	199
6.2.2	Synthesis of 2-(4-Chlorophenyl)-1,2-propanediol	201
6.2.3	Synthesis of 2-(4-Chlorophenyl)-1,2-hydroxypropanoic acid	202
	Conclusion	204
	References	204
6.3	Catalytic enantioselective epoxidation of <i>trans</i> -disubstituted and trisubstituted alkenes with arabinose-derived ulose	204
	<i>Tony K. M. Shing, Gulice Y.C. Leung and To Luk</i>	
6.3.1	Synthesis of 2',3'-Diisobutyl acetal	205
6.3.2	Synthesis of ulose	206
6.3.3	Asymmetric epoxidation of <i>trans</i> - α -Methylstilbene using ulose as catalyst at 0°C	208
	Conclusion	209
	References	210
6.4	VO(acac) ₂ /TBHP catalyzed epoxidation of 2-(2-Alkenyl)phenols. highly regio- and diastereoselective oxidative cyclisation to 2,3-Dihydrobenzofuranols and 3-Chromanols	211
	<i>Alessandra Lattanzi and Arrigo Scettri</i>	
6.4.1	VO(acac) ₂ /TBHP catalyzed epoxidation of 2-(3,7-Dimethyl-octa-2,6-dienyl)-phenol	212
6.4.2	VO(acac) ₂ /TBHP/TFA catalyzed oxidative cyclization of 2-(3,7-Dimethyl-octa-2,6-dienyl)-phenol	213
	Conclusion	214
	References	214

6.5	An Oxalolidinone ketone catalyst for the asymmetric epoxidation of <i>cis</i> -olefins.	215
	<i>David Goettel and Yan Shi</i>	
6.5.1	Amadori rearrangement to give 1-Dibenzylamino-1-deoxy-D-fructose	215
6.5.2	Acetal protection of 1-Dibenzylamino-1-deoxy-D-fructose	216
6.5.3	Hydrogenation of the Dibenzylamine.	217
6.5.4	Phosgene cyclization of aminoalcohol	218
6.5.5	Alcohol oxidation	220
6.5.6	Synthesis of ketone 2	221
6.5.7	Asymmetric epoxidation of <i>cis</i> - β -Methylstyrene	222
	Conclusion.	223
	References	224
6.6	α -Fluorotropinone immobilised on silica: a new stereoselective heterogeneous catalyst for epoxidation of alkenes with oxone	225
	<i>Giovanni Sartori, Alan Armstrong, Raimondo Maggi, Alessandro Mazzacani, Raffaella Sartorio, France Bigi and Belen Dominguez-Fernandez</i>	
6.6.1	Synthesis of silica KG-60-supported enantiomerically enriched α -Fluorotropinone.	225
6.6.2	Synthesis of enantiomerically enriched epoxides.	226
	Conclusion.	227
	References	228
6.7	Asymmetric epoxidation catalyzed by novel azacrown ether-type chiral quaternary ammonium salts under phase-transfer catalytic conditions.	228
	<i>Kazushige Hori, Keita Tani and Yasuo Tohda</i>	
6.7.1	Synthesis of precursor of the azacrown ether	229
6.7.2	Synthesis of the azacrown ether	230
6.7.3	Synthesis of the azacrown ether-type quaternary ammonium salt	232
6.7.4	Asymmetric epoxidation of (<i>E</i>)-Chalcone catalyzed by the azacrown ether-type quaternary ammonium salt as chiral PTC	233
	Conclusion.	234
	References	235
6.8	Enantioselective epoxidation of olefins using phase transfer conditions and a chiral [azepinium][TRISPHAT] salt as catalyst	235
	<i>Jérôme Vachon, Céline Pérollier, Alexandre Martinez and Jérôme Lacour</i>	
6.8.1	Enantioselective epoxidation of 1-Phenyl-3,4-dihydronaphthalene.	236
	Conclusion.	238
	References	239
6.9	Catalytic asymmetric epoxidation of α,β -unsaturated esters promoted by a Yttrium-biphenyldiol complex	239
	<i>Masakatsu Shibasaki, Hiroyuki Kakei and Shigeki Matsunaga</i>	
6.9.1	Synthesis of (<i>aS,R</i>)-6,6'-[(Propylene)dioxy]biphenyl-2,2'-diol	240
6.9.2	Synthesis of (<i>aS,R</i>)-2,2'-[Oxybis(ethylene)dioxy]-6,6'-[(propylene)dioxy]biphenyl	242
6.9.3	Synthesis of (<i>S</i>)-6,6'-[Oxybis(ethylene)dioxy]biphenyl-2,2'-diol	243
6.9.4	Enantiomeric enrichment of (<i>S</i>)-6,6'-[Oxybis(ethylene)dioxy]biphenyl-2,2'-diol	244
6.9.5	Catalytic asymmetric epoxidation of α,β -unsaturated esters	246
	References	248

6.10	Catalytic enantioselective epoxidation of α,β -enones with a binol-zinc-complex	249
	<i>Ana Minatti and Karl Heinz Dötz</i>	
6.10.1	Synthesis of (<i>E</i>)-(2 <i>S</i> ,3 <i>R</i>)-Phenyl-(3-phenyloxiran-2-yl)methanone.	249
	Conclusion.	250
	References.	251
6.11	Asymmetric epoxidation of Phenyl-2-(3'-pyridylvinyl)sulfone using poly-leucine hydrogen peroxide gel	251
	<i>Mike R. Pitts and John Whittall</i>	
6.11.1	Preparation of poly-leucine-hydrogen peroxide gel	252
6.11.2	Synthesis of Phenyl-2-(3'-pyridylvinyl) sulfone (2).	252
	References.	254
7	Oxidation of Ketones to Lactones or Enones	255
7.1	Synthesis of 2-(Phosphinophenyl)pyridine ligand and its application to palladium-catalyzed asymmetric Baeyer–Villiger oxidation of prochiral cyclobutanones	256
	<i>Katsuji Ito* and Tsutomu Katsuki</i>	
7.1.1	Synthesis of (7 <i>R</i>)-2-(2-Hydroxyphenyl)-7-isopropyl-6,7-dihydro-5 <i>H</i> -1-pyridine	256
7.1.2	2-[2-(Diphenylphosphinoyl)phenyl]-7-isopropyl-6,7-dihydro-5 <i>H</i> -1-pyridine	258
7.1.3	2-[2-(Diphenylphosphanyl)phenyl]-7-isopropyl-6,7-dihydro-5 <i>H</i> -1-pyridine	259
7.1.4	Asymmetric Baeyer–Villiger oxidation of 3-Phenylcyclobutanone	261
	Conclusion.	262
	References.	263
7.2	(<i>D</i>)-Codeinone from (<i>D</i>)-Dihydrocodeinone via the use of modified <i>o</i> -iodoxybenzoic acid (IBX). A convenient oxidation of ketones to enones	263
	<i>Paul Mather and John Whittall</i>	
7.2.1	Synthesis of IBX.	264
7.2.2	Synthesis of codeinone	264
	References.	266
8	Oxidative C–C Coupling	267
8.1	Enantioselective oxidative coupling of 2-Naphthols catalyzed by a novel chiral vanadium complex	267
	<i>Nan-Sheng Xie, Quan-Zhong Liu, Zhi-Bin Luo, Liu-Zhu Gong, Ai-Qiao Mi and Yao-Zhong Jiang</i>	
8.1.1	Synthesis of 3,3-Diformyl-2,2'-biphenol.	268
8.1.2	Synthesis of chiral vanadium complexes	270
8.1.3	Catalytic oxidative coupling of 7-Alkoxy-1-naphthols by chiral vanadium complexes	271
	Reference	272
8.2	Catalytic oxidative cross-coupling reaction of 2-Naphthol derivatives.	273
	<i>Shigeki Habaue and Tomohisa Temma</i>	
8.2.1	Synthesis of Methyl 2,2'-dihydroxy-1,1'-binaphthalene-3-carboxylate.	273

	Conclusion	274
	References	275
8.3	Oxidative coupling of benzenes with α,β -unsaturated aldehydes by Pd(OAc) ₂ /HPMoV/ O ₂ system	275
	<i>Tomoyuki Yamada, Satoshi Sakaguchi and Yasutaka Ishii</i>	
	8.3.1 Synthesis of Cinnamaldehyde	276
	Conclusion	278
	References	278
9	Oxidation of Sulfides and Sulfoxides	279
9.1	The first example of direct oxidation of sulfides to sulfones by an osmate-molecular oxygen system	280
	<i>Boyapati M. Choudary, Chinta Reddy, V. Reddy, Billakanti V. Prakash, Mannepalli L. Kantam and B. Sreedhar</i>	
	9.1.1 Synthesis of osmate exchanged Mg-Al layered double hydroxides (LDH-OsO ₄)	280
	9.1.2 Synthesis of Methyl phenyl sulfone or Methylsulfonylbenzene	281
	Conclusion	282
	References	283
9.2	Selective oxidation of sulfides to sulfoxides and sulfones using hydrogen peroxide in the presence of zirconium tetrachloride	283
	<i>Kiumar Bahrami</i>	
	9.2.1 Oxidation of Benzyl 4-bromobenzyl sulfide to Benzyl 4-bromobenzyl sulfoxide using H ₂ O ₂ in the presence of zirconium tetrachloride	284
	9.2.2 Oxidation of Benzyl 4-bromobenzyl sulfide to Benzyl 4-bromobenzyl sulfone using H ₂ O ₂ in the presence of zirconium tetrachloride	286
	Conclusion	287
	References	287
9.3	WO ₃ -30 % H ₂ O ₂ -cinchona alkaloids: a new heterogeneous catalytic system for asymmetric oxidation and kinetic resolution of racemic sulfoxides	288
	<i>Vinay V. Thakur and A. Sudalai</i>	
	9.3.1 Synthesis of (<i>R</i>)-2-[[[3-Methyl-4-(2,2,2-trifluoroethoxy)-2- pyridyl]methyl]sulfinyl]-1 <i>H</i> -benzimidazole {(<i>R</i>)-(+)-Lansoprazole}	288
	9.3.2 Synthesis of (<i>R</i>)-(+)-Phenyl benzyl sulfoxide	290
	Conclusion	293
	References	293
9.4	Benzyl-4,6- <i>O</i> -isopropylidene- α -D-glucopyranoside, 2-deoxy- 2-[[[2-hydroxy-3,5-di- <i>tert</i> -butylphenyl]methylene]amino] as a ligand for vanadium-catalyzed asymmetric oxidation of sulfides	293
	<i>Raffaella Del Litto, Guiseppina Roviello and Francesco Ruffo</i>	
	9.4.1 Synthesis of Benzyl-4,6- <i>O</i> -isopropylidene- α -D-glucopyranoside, 2-deoxy-2-[[[2-hydroxy-3,5-di- <i>tert</i> -butylphenyl]methylene]imine]	294
	9.4.2 Oxidation of Thioanisole	295
	Conclusion	296
	References	296
9.5	Asymmetric sulfoxidation of aryl methyl sulfides with hydrogen peroxide in water	297
	<i>Alessandro Scarso and Giorgio Strukul</i>	

CONTENTS

xv

9.5.1	Synthesis of complex $(R)\text{-BINAP)PtCl}_2$	298
9.5.2	Synthesis of complex $[(R)\text{-BINAP)Pt}(\text{OH})_2(\text{BF}_4)_2$	299
9.5.3	Stereoselective catalytic oxidation of aryl methyl sulfides	300
	Conclusion	300
	References	301
	Index	303