

Plant Biology

Alison M. Smith
George Coupland
Liam Dolan
Nicholas Harberd
Jonathan Jones
Cathie Martin
Robert Sablowski
Abigail Amey

CONTENTS

Preface	v		
Acknowledgments	vi		
Contents in Brief	vii		
Chapter 1 Origins	1		
1.1 Earth, Cells, and Photosynthesis	2		
The earth formed 4.6 billion years ago	2		
Photosynthesis evolved by 3.8 billion years ago	4		
Oxygen-producing photosynthesis was widespread by 2.2 billion years ago	5		
Photosynthetic cyanobacteria produced an oxygen-rich atmosphere	6		
Early life on earth evolved in the absence of a protective atmospheric ozone layer	6		
1.2 Eukaryotic Cells	7		
Photosynthetic eukaryotic cells arose from two endosymbiotic events	7		
Several groups of photosynthetic organisms are derived from the endosymbiotic event that gave rise to plastids	8		
Fossil evidence indicates that eukaryotic organisms had evolved by 2.7 billion years ago and multicellular organisms by 1.25 billion years ago	10		
Animals and algae diversified in the Early Cambrian Period	11		
Box 1–1 What DNA Can Reveal about Phylogeny and Evolution	14		
1.3 Land Plants	12		
Green plants are monophyletic	13		
Land plants may be descended from plants related to charophycean (charophyte) algae	13		
Microfossils indicate that the first land plants appeared in the Middle Ordovician Period, about 475 million years ago	15		
Plant diversity increased in the Silurian and Devonian Periods	16		
The number of sporangia distinguishes the first land plants from their evolutionary descendants	16		
		Increases in plant size were accompanied by evolution of a vascular system	18
		Some of the earliest vascular plants were related to extant lycophytes	19
		Horsetails, ferns, and seed plants are derived from a leafless group of plants of the Early Devonian Period, 400 million years ago	20
		Ferns and horsetails evolved in the Devonian Period	22
		Chemical and cellular complexity increased early in the evolution of land plants	23
		Atmospheric CO ₂ and O ₂ levels are determined by rates of photosynthesis and carbon burial	24
		The evolution of land plants was at least partly responsible for the decrease in atmospheric CO ₂ beginning 450 million years ago	25
		The mid-Paleozoic decrease in atmospheric CO ₂ was a driving force in the evolution of big leaves	26
		1.4 Seed Plants	27
		Seeds contain the genetic products of fertilization protected by tissue derived from the sporophyte	28
		Seed plants evolved in the Devonian and diversified in the Permian, 290 to 250 million years ago	29
		The sporophyte phase became dominant in the land-plant life cycle in the Devonian Period	30
		Five groups of seed plants live on earth today	33
		1.5 Angiosperms	33
		Angiosperms appear in the fossil record in the Early Cretaceous Period, about 135 million years ago	34
		Angiosperms evolved in the tropics and then spread to higher latitudes	34
		<i>Amborella trichopoda</i> is sister to all living angiosperms	35
		Eudicots are distinguished from other flowering plants by the number of pollen apertures	37
		The earliest angiosperm flowers were small with many parts	38
		Monocots are a monophyletic group	38
		The grass family (Poaceae) evolved about 60 million years ago but diversified more recently	39

Chapter 2 Genomes	43		
2.1 The Nuclear Genome: Chromosomes	44		
		Genome sequencing allows the development of methods to monitor the activity of many genes simultaneously	78
2.2 Chromosomal DNA	45	2.6 Cytoplasmic Genomes	79
Specialized, repetitive DNA sequences are found in the centromeres and telomeres	45	Plastids and mitochondria evolved from bacteria engulfed by other cells	80
Nuclear genes are transcribed into several types of RNA	47	Organellar genes do not follow Mendel's laws of inheritance	80
Plant chromosomes contain many mobile genetic elements	49	The genomes of plastids and mitochondria have been reduced during evolution	80
2.3 Nuclear Gene Regulation	52	Most polypeptides in organelles are encoded by the nuclear genome and targeted to the organelles	81
Regulatory sequences and transcription factors control where and when a gene is transcribed	52	Replication and recombination of plastid DNA is not tightly coupled to cell division	82
Gene activity can be regulated by chemical changes in the DNA and proteins of chromatin	57	Gene expression has common features in plastids and eubacteria	82
Chromatin modification can be inherited through cell division	59	Plastids contain two distinct RNA polymerases	83
Gene function is also controlled at the RNA level	60	Post-transcriptional processes are important in regulating plastid gene expression	84
Small regulatory RNAs control mRNA function	61	Organellar transcripts undergo RNA editing	85
Small RNAs can direct chromatin modification to specific DNA sequences	63	Post-translational processes contribute to maintaining the correct ratio of nuclear- and plastid-encoded components of multisubunit complexes	85
Box 2-1 Transcription Factors: Combinatorial Control	53	Developmental regulation of plastid gene expression includes signaling pathways between plastids and the nucleus	86
2.4 Genome Sequences	64		
The Arabidopsis genome was the first plant genome to be fully sequenced	65	Chapter 3 Cells	91
Genome sequences are analyzed to identify individual genes	65	3.1 The Cell Cycle	93
Sequencing of the Arabidopsis genome revealed a complexity similar to that of animal genomes and a large proportion of plant-specific genes	66	Transition from one phase of the cell cycle to the next is regulated by a complex set of mechanisms	93
Comparisons among plant genomes reveal conserved and divergent features	68	The cell cycle in plants is controlled by developmental and environmental inputs	98
Most angiosperms have undergone genome duplication during their evolution	68	Many differentiating cells undergo endoreduplication: DNA replication without nuclear and cell division	99
Genes can acquire new functions by duplication and divergence	70	Box 3-1 The Nucleus	95
The order of genes is conserved between closely related plant species	72	3.2 Cell Division	102
2.5 Genomes and Biotechnology	73	The cytoskeleton moves cellular components during cell division	102
Mutated genes can be localized on the genome by co-segregation with known markers	74	A preprophase band forms at the site of the future cell wall	104
Genes that are mutated by insertion of DNA can be isolated by detecting the inserted sequence	74	Replicated pairs of chromosomes are separated on a spindle of microtubules	105
Genes can be screened for mutations at the DNA level independent of phenotype	75	Microtubules direct the formation of the phragmoplast, which orchestrates deposition of the new cell wall	106
RNA interference is an alternative method to knock out gene function	76	Vesicles carry material from the Golgi apparatus to the newly forming cell wall	109
Multigenic inheritance is analyzed by mapping quantitative trait loci (QTLs)	77	Meiosis is a specialized type of cell division that gives rise to haploid cells and genetic variation	112
		Box 3-2 The Cytoskeleton	103

3.3 Organelles	116	Chapter 4 Metabolism	167
Plastids and mitochondria replicate independent of cell division	117	4.1 Control of Metabolic Pathways	169
Plastid and mitochondrial biogenesis involves post-translational import of many proteins	119	Compartmentation increases the potential for metabolic diversity	169
The endomembrane system delivers proteins to the cell surface and to vacuoles	122	Metabolic processes are coordinated and controlled by regulation of enzyme activities	170
Organelles move around the cell on actin filaments	127	4.2 Carbon Assimilation: Photosynthesis	174
3.4 Primary Cell Wall	128	Net carbon assimilation occurs in the Calvin cycle	175
The matrix of the cell wall consists of pectins and hemicelluloses	129	Energy for carbon assimilation is generated by light-harvesting processes in the chloroplast thylakoids	175
Cellulose is synthesized at the cell surface after the cell plate has formed	130	Light energy is captured by chlorophyll molecules and transferred to reaction centers	177
Carbohydrate components of the cell wall interact to form a strong and flexible structure	132	Electron transfer between two reaction centers via an electron transport chain reduces NADP ⁺ and generates a proton gradient across the thylakoid membrane	180
Glycoproteins and enzymes have important functions in the cell wall	134	The proton gradient drives the synthesis of ATP by an ATP synthase complex	184
Plasmodesmata form channels between cells	135	Light-harvesting processes are regulated to maximize the dissipation of excess excitation energy	186
3.5 Cell Expansion and Cell Shape	138	Carbon assimilation and energy supply are coordinated by complex regulation of Calvin cycle enzymes	188
The properties of the plasma membrane determine the composition of the cell and mediate its interactions with the environment	138	Sucrose synthesis is tightly controlled by the rate of photosynthesis and the demand for carbon by nonphotosynthetic parts of the plant	190
Proton transport across the plasma membrane generates electrical and proton gradients that drive other transport processes	138	Synthesis of starch allows the photosynthetic rate to remain high when sucrose synthesis is restricted	195
Movement of water across the plasma membrane is facilitated by aquaporins	141	Box 4–1 Light	178
Movement of solutes into the cell vacuole drives cell expansion	142	4.3 Photorespiration	197
The vacuole acts as a storage and sequestration compartment	144	Rubisco can use oxygen instead of carbon dioxide as substrate	197
Coordinated ion transport and water movement drive stomatal opening	146	Photorespiratory metabolism has implications for both the carbon and the nitrogen economy of the leaf	200
The direction of cell expansion is determined by microtubules in the cell cortex	149	C4 plants eliminate photorespiration by a mechanism that concentrates carbon dioxide	203
Actin filaments direct new material to the cell surface during cell expansion	151	4.4 Sucrose Transport	210
In root hair cells and pollen tubes, cell expansion is localized to the cell tips	152	Sucrose moves to nonphotosynthetic parts of the plant via the phloem	210
3.6 Secondary Cell Wall and Cuticle	154	Phloem loading may be apoplastic or symplastic	210
The structure and components of the secondary cell wall vary from one cell type to another	154	The path of sucrose unloading from the phloem depends on the type of plant organ	215
Lignin is a major component of many secondary cell walls	155	The supply of assimilate from the leaf is coordinated with demand elsewhere in the plant	216
Lignification is a defining characteristic of xylem vessels and tracheids	160	4.5 Nonphotosynthetic Generation of Energy and Precursors	217
Wood is formed by secondary growth of vascular tissues	160	Interconversion of sucrose and hexose phosphates allows sensitive regulation of sucrose metabolism	218
The cuticle forms a hydrophobic barrier on the aerial parts of the plant	163		

Glycolysis and the oxidative pentose phosphate pathway generate reducing power, ATP, and precursors for biosynthetic pathways	220		
The Krebs cycle and mitochondrial electron transport chains provide the main source of ATP in nonphotosynthesizing cells	222		
Partitioning of sucrose among “metabolic backbone” pathways is extremely flexible and is related to cell function	230		
4.6 Carbon Storage	233		
Sucrose is stored in the vacuole	234		
The starch granule is a semi-crystalline structure synthesized by small families of starch synthases and starch-branching enzymes	235		
The pathway of starch degradation depends on the type of plant organ	239		
Some plants store soluble fructose polymers rather than starch	242		
Storage lipids are synthesized from fatty acids in the endoplasmic reticulum	242		
The fatty acid composition of storage lipids varies among species	244		
Triacylglycerols are converted to sugars by β oxidation and gluconeogenesis	249		
Sugars may act as signals that determine the extent of carbon storage	251		
4.7 Plastid Metabolism	253		
Plastids exchange specific metabolites with the cytosol via <i>metabolite transporters</i>	253		
Fatty acids are synthesized by an enzyme complex in plastids	256		
Membrane lipid synthesis in plastids proceeds via a “prokaryotic” pathway different from the “eukaryotic” pathway elsewhere in the cell	259		
Different pathways of terpenoid synthesis in the plastid and the cytosol give rise to different products	262		
Tetrapyrroles, the precursors of chlorophyll and heme, are synthesized in plastids	265		
4.8 Nitrogen Assimilation	270		
Plants contain several types of nitrate transporter, regulated in response to different signals	270		
Nitrate reductase activity is regulated at many different levels	272		
Assimilation of nitrogen into amino acids is coupled to demand, nitrate availability, and availability of biosynthetic precursors	274		
Amino acid biosynthesis is partly controlled by feedback regulation	276		
Nitrogen is stored as amino acids and specific storage proteins	281		
		4.9 Phosphorus, Sulfur, and Iron Assimilation	284
		The availability of phosphorus is a major limitation on plant growth	287
		Sulfur is taken up as sulfate, then reduced to sulfide and assimilated into cysteine	288
		Iron uptake requires specialized mechanisms to increase iron solubility in the soil	291
		4.10 Movement of Water and Minerals	293
		Water moves from the soil to the leaves, where it is lost in transpiration	293
		Water moves from roots to leaves by a hydraulic mechanism	294
		The movement of mineral nutrients in the plant involves both xylem and phloem	296
		Chapter 5 Development	301
		5.1 Overview of Plant Development	302
		Multicellularity evolved independently in plants and animals	304
		<i>Volvox</i> is a simple system in which to study the genetic basis of multicellularity	305
		5.2 Embryo and Seed Development	306
		External cues establish the apical-basal axis in the <i>Fucus</i> embryo	307
		The cell wall directs the fate of cells in the <i>Fucus</i> embryo	308
		Embryo development in higher plants occurs within the seed	309
		The fate of embryonic cells is defined by their position	310
		Progressive polarization of auxin transporters mediates formation of the basal pole in embryos	312
		Radial cell pattern in the embryonic root and hypocotyl is defined by the SCARECROW and SHORT ROOT transcription factors	313
		Clues from apical-basal and radial patterning of the embryo are combined to position the root meristem	314
		The shoot meristem is established gradually and independent of the root meristem	315
		The endosperm and embryo develop in parallel	316
		Division of the cells that give rise to endosperm is repressed until fertilization	317
		After embryo and endosperm development, seeds usually become dormant	319
		Box 5–1 Clonal Analysis	311
		5.3 Root Development	320
		Plant roots evolved independently at least twice	321
		Roots have several zones containing cells at successive stages of differentiation	321

The <i>Arabidopsis</i> root has simple cellular organization	322	In the ABC model of floral organ identity, each type of organ is determined by a specific combination of homeotic genes	354
Cell fate depends on the cell's position in the root	323	Floral organ identity genes are conserved throughout the angiosperms	357
Genetic analysis confirms the position-dependent specification of cell type	324	Asymmetrical growth of floral organs gives rise to bilaterally symmetrical flowers	358
Lateral root development requires auxin	325	Additional regulatory genes control later stages of floral organ development	358
Box 5-2 Stem Cells in Plants and Animals	323		
5.4 Shoot Development	326	5.6 From Sporophyte to Gametophyte	360
Cells in the shoot apical meristem are organized in radial zones and in concentric layers	327	The male gametophyte is the pollen grain, with a vegetative cell, gametes, and a tough cell wall	360
The number of new meristem cells is constantly balanced by the number that form new organs	330	Pollen development is aided by the surrounding sporophyte tissues	362
Organ primordia emerge from the flanks of the meristem in a repetitive pattern	332	The female gametophyte develops in the ovule, which contains gametes for the two fertilization events that form the zygote and the endosperm	364
Changes in gene expression precede primordium emergence	333	Development of the female gametophyte is coordinated with development of the sporophyte tissues of the ovule	365
Development of compound leaves is associated with expression of meristem genes during early leaf development	334	A pollen grain germinates on the carpel and forms a tube that transports the sperm nuclei toward the ovule	365
Leaves are shaped by organized cell division followed by a period of cell expansion and differentiation	335	Growth of the pollen tube is oriented by long-range signals in the carpel tissues and short-range signals produced by the ovule	366
Different regions of the leaf primordium acquire different fates early in development	335	Plants have mechanisms that allow the growth only of pollen tubes carrying specific genes	367
Specific genes regulate the differences between the two faces of the leaf	337	<i>Self-incompatibility</i> can be gametophytic or sporophytic, depending on the origin of the pollen protein recognized	367
Lateral growth requires the boundary between the dorsal and ventral sides of the leaf	338	Angiosperms have double fertilization	369
The leaf reaches its final shape and size by regulated cell division and cell expansion	339	Genes from the male and female gametes are not expressed equally after fertilization	370
Leaf growth is accompanied by development of an increasingly elaborate vascular system, which is controlled by auxin transport	340	Some plants can form seeds without fertilization	371
Cell communication and oriented cell divisions control the placement of specialized cell types in the leaf	342		
Leaf senescence is an active process that retrieves nutrients from leaves at the end of their useful lifespan	344	Chapter 6 Environmental Signals	377
Branches originate from lateral meristems whose growth is influenced by the apical meristem	346	6.1 Seed Germination	378
Internodes grow by cell division and cell elongation, controlled by gibberellins	347	6.2 Light and Photoreceptors	380
A layer of meristem cells generates vascular tissues and causes secondary thickening of the stem	349	Plant development proceeds along distinct pathways in light and dark	380
5.5 From Vegetative to Reproductive Development	350	Distinct photoreceptors detect light of different wavelengths	381
Reproductive structures in angiosperms are produced by floral and inflorescence meristems	350	Phytochromes are converted from an inactive to an active form by exposure to red light	382
Development of floral meristems is initiated by a conserved regulatory gene	351	Distinct forms of phytochrome have different functions	385
The expression pattern of LEAFY-like genes determines inflorescence architecture	351	Phytochrome plays a role in shade avoidance	387
Flowers vary greatly in appearance, but their basic structure is directed by highly conserved genes	353	Cryptochromes are blue-light receptors with specific and overlapping functions	388
		Phototropins are blue-light receptors involved in phototropism, stomatal opening, and chloroplast migration	390

Some photoreceptors respond to red and blue light	392	Statoliths are key to graviperception in stems, hypocotyls, and roots	430
Biochemical and genetic studies provide information on the components of the phytochrome signal-transduction pathway	392	Columella cells of the root cap are the site of graviperception in the growing root	430
6.3 Seedling Development	395	The endodermal cell layer is the site of graviperception in growing stems and hypocotyls	431
Ethylene is synthesized from methionine in a pathway controlled by a family of genes	396	Mutations in auxin signaling or transport cause defects in root gravitropism	431
Genetic analysis has identified components of the ethylene signal-transduction pathway	396	The extent of lateral root elongation varies in response to soil nutrient levels	432
The ethylene response is negatively regulated by binding of ethylene to its receptors	398	Chapter 7 Environmental Stress	437
Inactivation of CTR1 allows activation of downstream components of the ethylene signaling chain	399	7.1 Light as Stress	438
Ethylene interacts with other signaling pathways	400	Photosystem II is highly sensitive to too much light	439
The light responses of seedlings are repressed in the dark	400	High light induces nonphotochemical quenching, a short-term protective mechanism against photooxidation	439
COP1 and the COP9 signalosome function by destabilizing proteins required for photomorphogenesis	402	Vitamin E-type antioxidants also protect PSII under light stress	443
Brassinosteroids are required for repression of photomorphogenesis in the dark and other important functions in plant development	403	Photodamage to photosystem II is quickly repaired in light stress-tolerant plants	444
6.4 Flowering	407	Some plants, such as winter evergreens, have mechanisms for longer-term protection against light stress	445
Reproductive development in many plants is controlled by photoperiod	408	Low light leads to changes in leaf architecture, chloroplast structure and orientation, and life cycle	447
Phytochromes and cryptochromes act as light receptors in the photoperiodic control of flowering	410	Ultraviolet irradiation damages DNA and proteins	449
Circadian rhythms control the expression of many plant genes and affect the photoperiodic control of flowering	411	Resistance to UV light involves the production of specialized plant metabolites, as well as morphological changes	451
Circadian rhythms in plants result from input of environmental signals, a central oscillator, and output of rhythmic responses	413	7.2 High Temperature	452
Substances produced in leaves can promote or inhibit flowering	416	High temperature induces the production of heat shock proteins	453
Similar groups of genes are involved in photoperiodic control of flowering in <i>Arabidopsis</i> and in rice	418	Molecular chaperones ensure the correct folding of proteins under all conditions	454
Vernalization is detected in the apex and controls flowering time in many plants	422	Families of heat shock proteins play different roles in the heat stress response in different species	454
Genetic variation in the control of flowering may be important in the adaptation of plants to different environments	424	Synthesis of heat shock proteins is controlled at the transcriptional level	455
Vernalization response in <i>Arabidopsis</i> involves modification of histones at the <i>FLC</i> gene, which is also regulated by the autonomous flowering pathway	426	Some plants have developmental adaptations to heat stress	456
Photoperiodic and vernalization pathways of <i>Arabidopsis</i> converge to regulate the transcription of a small set of floral integrator genes	428	7.3 Water Deficit	456
6.5 Root and Shoot Growth	429	Water deficit occurs as a result of drought, salinity, and low temperature	456
Plant growth is affected by gravitational stimuli	429	Plants use abscisic acid as a signal to induce responses to water deficit	457
		Plants also use ABA-independent signaling pathways to respond to drought	460
		Abscisic acid regulates stomatal opening to control water loss	461
		Drought-induced proteins synthesize and transport osmolytes	462

Ion channels and aquaporins are regulated in response to water stress	464	7.7 Oxidative Stress	493
Many plant species adopt metabolic specialization under drought stress	465	Reactive oxygen species are produced during normal metabolism, but also accumulate under a range of environmental stress conditions	493
Plants that tolerate extreme desiccation have a modified sugar metabolism	467	Ascorbate metabolism is central to the elimination of reactive oxygen species	494
Many plant species adapted to arid conditions have specialized morphology	468	Hydrogen peroxide signals oxidative stress	496
Rapid life cycling during water availability is common in plants of arid regions	471	Ascorbate metabolism is central to responses to oxidative stress	496
7.4 Salt Stress	472	Chapter 8 Interactions with Other Organisms	499
Salt stress disrupts homeostasis in water potential and ion distribution	472	8.1 Microbial Pathogens	501
Salt stress is signaled by ABA-dependent and ABA-independent pathways	472	Most pathogens can be classified as biotrophs or necrotrophs	502
Adaptations to salt stress principally involve internal sequestration of salts	473	Pathogens enter plants via several different routes	503
Physiological adaptations to salt stress include modulation of guard cell function	476	Pathogen infections lead to a broad range of disease symptoms	506
Morphological adaptations to salt stress include salt-secreting trichomes and bladders	476	Many pathogens produce effector molecules that influence their interactions with the host plant	507
Osmotic stress stimulates reproduction in some halophytes	479	<i>Agrobacterium</i> transfers its DNA (T-DNA) into plant cells to modify plant growth and feed the bacterium, and this transfer system is used in biotechnology	511
7.5 Cold	479	Some pathogen effector molecules are recognized by the plant and trigger defense mechanisms	515
Low temperature is similar to water deficit as an environmental stress	479	The products of some bacterial <i>avr</i> genes act inside the plant cell	516
Temperate plants acclimated by prior exposure to low temperatures are resistant to freezing damage	480	The functions of fungal and oomycete effector molecules are poorly understood	518
Exposure to low temperatures induces cold-regulated (<i>COR</i>) genes	481	8.2 Pests and Parasites	519
Expression of the transcriptional activator CBF1 induces <i>COR</i> gene expression and cold tolerance	481	Parasitic nematodes form intimate associations with host plants	519
Low-temperature signaling involves increases in intracellular calcium	483	Insects cause extensive losses in crop plants, both directly and by facilitating infection by pathogens	521
ABA-dependent and ABA-independent pathways signal in response to cold	483	Some plants are plant pathogens	522
Plant species of warm climates are chill-sensitive	483	8.3 Viruses and Viroids	524
Vernalization and cold acclimation are closely linked processes in wheat and other cereal crops	483	Viruses and viroids are a diverse and sophisticated set of parasites	524
7.6 Anaerobic Stress	484	Different types of plant viruses have different structures and replication mechanisms	525
Water-logging is a cause of hypoxic or anoxic stress for plants	485	8.4 Defenses	529
Hypoxia is signaled by a Rop-mediated signaling pathway involving transient induction of ROS	485	Basal defense mechanisms are triggered by pathogen-associated molecular patterns (PAMPs)	530
Anoxia induces shifts in primary metabolism	486	R proteins and many other plant proteins involved in defense carry leucine-rich repeats	534
Aerenchyma facilitates long-distance oxygen transport in flood-tolerant plants	488	R genes encode families of proteins involved in recognition and signal transduction	535
Water-logging is associated with other developmental adaptations that increase plant survival	490	Most R proteins do not directly recognize pathogen effector molecules	536
Plants synthesize oxygen-binding proteins under hypoxic conditions	493	R gene polymorphism restricts disease in natural populations	538

<i>R</i> genes have been selected in crop breeding from the earliest times	540	9.2 Scientific Plant Breeding	583
Insensitivity to toxins is important in plant defense against necrotrophs	541	Scientific approaches to crop plant improvement have resulted in substantial changes in the genetic structure of many crops	583
Plants synthesize antibiotic compounds that confer resistance to some microbes and herbivores	542	Triticale is a synthetic domesticated crop species	585
Disease resistance is often associated with the localized death of plant cells	548	Disease resistance is an important determinant of yield and can be addressed both by plant breeding and by crop management	586
In systemic resistance, plants are “immunized” by biological challenges that lead to cell death	549	Mutations in genes affecting fruit color, fruit ripening, and fruit drop have been used in tomato breeding programs	587
Wounding and insect feeding induce complex plant defense mechanisms	552	In the “Green Revolution,” the use of dwarfing mutations of wheat and rice resulted in major increases in crop yield	588
Chewing insects provoke release of volatile compounds that attract other insects	554	Heterosis also results in major increases in crop yields	590
RNA silencing is important in plant resistance to viruses	555	Cytoplasmic male sterility facilitates the production of F1 hybrids	592
8.5 Cooperation	557	9.3 Biotechnology	593
Many plant species are pollinated by animals	557	<i>Agrobacterium</i> -mediated gene transfer is a widely used method for generating transgenic plants	593
Symbiotic nitrogen fixation involves specialized interactions of plants and bacteria	560	Particle bombardment-mediated gene transfer is an alternative means of generating transgenic plants	594
Mycorrhizal fungi form intimate symbioses with plant roots	568	Herbicide resistance in transgenic crops facilitates weed control	595
Chapter 9 Domestication and Agriculture	573	Transgenic expression of <i>Bacillus thuringiensis</i> (Bt) crystal protein in crop plants confers insect resistance and increased yield	596
9.1 Domestication	574	Many different crop plant traits can potentially be improved by transgenesis	596
The domestication of crop species involved selection by humans	574	“The future is green”: The relationship between plants and people will continue to develop	599
The difference between maize and its wild ancestor, teosinte, can be explained by allelic variation at five different loci	576	Glossary	603
Alterations in the expression of the gene <i>teosinte branched</i> played an important role in the domestication of maize	578	Figure Acknowledgments	631
The <i>teosinte glume architecture</i> gene regulates glume size and hardness	579	Index	635
Cultivated wheat is polyploid	580		
Cauliflower arose through mutation of a meristem-identity gene	581		
Increase in fruit size occurred early in the domestication of tomato	583		