

Contents

| | |
|---|------------------|
| <i>List of contributors</i> | <i>page</i> xiii |
| <i>Preface</i> | xvi |
| Part I Principles and techniques | 1 |
| 1 General principles and characteristics of optical magnetometers | 3 |
| <i>D. F. Jackson Kimball, E. B. Alexandrov, and D. Budker</i> | |
| 1.1 Introduction | 3 |
| 1.1.1 Fundamental sensitivity limits | 4 |
| 1.1.2 Zeeman shifts and atomic spin precession | 5 |
| 1.1.3 Quantum beats and dynamic range | 8 |
| 1.2 Model of an optical magnetometer | 8 |
| 1.3 Density matrix and atomic polarization moments | 13 |
| 1.4 Sensitivity and accuracy | 16 |
| 1.4.1 Variational sensitivity (short-term resolution) and long-term stability | 16 |
| 1.4.2 Parameter optimization | 18 |
| 1.4.3 Absolute accuracy and systematic errors | 19 |
| 1.5 Vector and scalar magnetometers | 20 |
| 1.6 Applications | 21 |
| 2 Quantum noise in atomic magnetometers | 25 |
| <i>M. V. Romalis</i> | |
| 2.1 Introduction | 25 |
| 2.2 Spin-projection noise | 26 |
| 2.3 Faraday rotation measurements | 26 |
| 2.4 Quantum back-action | 27 |
| 2.5 Time correlation of spin-projection noise | 28 |
| 2.6 Conditions for spin-noise dominance | 30 |
| 2.7 Spin projection limits on magnetic field sensitivity | 32 |
| 2.8 Spin squeezing and atomic magnetometry | 36 |
| 2.9 Conclusion | 37 |

| | | |
|-------|--|----|
| 3 | Quantum noise, squeezing, and entanglement in radiofrequency optical magnetometers | 40 |
| | <i>K. Jensen and I. S. Polzik</i> | |
| 3.1 | Sources of noise | 40 |
| 3.1.1 | Atomic projection noise | 40 |
| 3.1.2 | Photon shot noise | 41 |
| 3.1.3 | Back-action noise and QND measurements | 42 |
| 3.1.4 | Technical (classical) noise | 42 |
| 3.1.5 | Entanglement and spin squeezing | 42 |
| 3.2 | A pulsed radiofrequency magnetometer and the projection noise limit | 43 |
| 3.2.1 | Pulsed RF magnetometry | 44 |
| 3.2.2 | Sensitivity and bandwidth | 45 |
| 3.3 | Light-atom interaction | 46 |
| 3.3.1 | A spin-polarized atomic ensemble interacting with polarized light | 47 |
| 3.3.2 | Conditional spin squeezing | 48 |
| 3.3.3 | Larmor precession, back-action noise, and two atomic ensembles | 48 |
| 3.3.4 | Swap and squeezing interaction | 49 |
| 3.4 | Demonstration of high-sensitivity, projection-noise-limited magnetometry | 50 |
| 3.4.1 | Setup, pulse sequence, and procedure | 50 |
| 3.4.2 | The projection-noise-limited magnetometer | 52 |
| 3.5 | Demonstration of entanglement-assisted magnetometry | 54 |
| 3.6 | Conclusions | 57 |
| 4 | M_x and M_z magnetometers | 60 |
| | <i>E. B. Alexandrov and A. K. Vershovskiy</i> | |
| 4.1 | Dynamics of magnetic resonance in an alternating field | 60 |
| 4.1.1 | Bloch equations and Bloch sphere | 60 |
| 4.1.2 | Types of magnetic resonance signals: M_z and M_x signals | 62 |
| 4.2 | M_z and M_x magnetometers: general principles | 63 |
| 4.2.1 | Advantages and disadvantages of M_z magnetometers | 66 |
| 4.2.2 | Advantages and disadvantages of M_x magnetometers | 67 |
| 4.2.3 | Attempts to combine advantages of M_x and M_z magnetometers: M_x - M_z tandems | 72 |
| 4.3 | Applications: radio-optical M_x and M_z magnetometers | 73 |
| 4.3.1 | Alkali M_z magnetometers | 73 |
| 4.3.2 | M_x magnetometers | 75 |
| 4.3.3 | M_x - M_z tandems | 79 |
| 4.4 | Summary: M_x and M_z scheme limitations, prospects, and application areas | 82 |

| | | |
|-------|---|-----|
| 5 | Spin-exchange-relaxation-free (SERF) magnetometers | 85 |
| | <i>I. Savukov and S. J. Seltzer</i> | |
| 5.1 | Introduction | 85 |
| 5.2 | Spin exchange collisions | 86 |
| 5.2.1 | The density-matrix equation | 86 |
| 5.2.2 | Simple model of spin exchange | 90 |
| 5.3 | Bloch equation description | 92 |
| 5.4 | Experimental realization | 95 |
| 5.4.1 | Classic SERF atomic magnetometer arrangement | 95 |
| 5.4.2 | Zeroing the magnetic field | 98 |
| 5.4.3 | Use of antirelaxation coatings | 98 |
| 5.4.4 | Comparison with SQUIDS | 99 |
| 5.5 | Fundamental sensitivity | 101 |
| 6 | Optical magnetometry with modulated light | 104 |
| | <i>D. F. Jackson Kimball, S. Pustelny, V. V. Yashchuk, and D. Budker</i> | |
| 6.1 | Introduction | 104 |
| 6.2 | Typical experimental arrangements | 106 |
| 6.3 | Resonances in the magnetic field dependence | 108 |
| 6.3.1 | Frequency modulation | 108 |
| 6.3.2 | Amplitude modulation | 111 |
| 6.3.3 | Polarization modulation | 113 |
| 6.4 | Effects at high light powers | 113 |
| 6.5 | Nonlinear Zeeman effect | 116 |
| 6.6 | Magnetometric measurements with modulated light | 118 |
| 6.7 | Conclusion | 122 |
| 7 | Microfabricated atomic magnetometers | 125 |
| | <i>S. Knappe and J. Kitching</i> | |
| 7.1 | Introduction | 125 |
| 7.2 | Sensitivity scaling with size | 126 |
| 7.3 | Sensor fabrication | 131 |
| 7.4 | Vapor cells | 133 |
| 7.5 | Heating and thermal management | 134 |
| 7.6 | Performance | 135 |
| 7.7 | Applications of microfabricated magnetometers | 137 |
| 7.8 | Outlook | 139 |
| 8 | Optical magnetometry with nitrogen-vacancy centers in diamond | 142 |
| | <i>V. M. Acosta, D. Budker, P. R. Hemmer, J. R. Maze, and R. L. Walsworth</i> | |
| 8.1 | Introduction | 142 |
| 8.1.1 | Comparison with existing technologies | 143 |
| 8.2 | Historical background | 144 |
| 8.2.1 | Single-spin optically detected magnetic resonance | 145 |
| 8.3 | NV center physics | 146 |

| | | |
|--------|--|-----|
| 8.3.1 | Intersystem crossing and optical pumping | 146 |
| 8.3.2 | Ground-state level structure and ODMR-based magnetometry | 148 |
| 8.3.3 | Interaction with environment | 150 |
| 8.4 | Experimental realizations | 152 |
| 8.4.1 | Near-field scanning probes and single-NV magnetometry | 152 |
| 8.4.2 | Wide-field array magnetic imaging | 157 |
| 8.4.3 | NV-ensemble magnetometers | 158 |
| 8.5 | Outlook | 161 |
| 9 | Magnetometry with cold atoms | 167 |
| | <i>W. Gawlik and J. M. Higbie</i> | |
| 9.1 | Introduction | 167 |
| 9.2 | Experimental conditions | 168 |
| 9.2.1 | Constraints and advantages of using cold atoms for magnetometry | 168 |
| 9.2.2 | Cold samples of atoms above quantum degeneracy | 168 |
| 9.3 | Linear Faraday rotation with trapped atoms | 170 |
| 9.4 | Nonlinear Faraday rotation | 173 |
| 9.4.1 | Low-field, DC magnetometry | 173 |
| 9.4.2 | Coherence evolution | 174 |
| 9.4.3 | High-field, amplitude-modulated magneto-optical rotation | 175 |
| 9.4.4 | Paramagnetic nonlinear rotation | 175 |
| 9.5 | Magnetometry with ultra-cold atoms | 176 |
| 9.5.1 | Overview of ultra-cold atomic magnetometry methods | 176 |
| 9.5.2 | Figures of merit | 180 |
| 9.5.3 | Details of spinor magnetometry | 182 |
| 9.5.4 | Comparison with thermal-atom magnetometry | 185 |
| 9.5.5 | Applications | 187 |
| 10 | Helium magnetometers | 190 |
| | <i>R. E. Slocum, D. D. McGregor, and A. W. Brown</i> | |
| 10.1 | Introduction | 190 |
| 10.2 | Helium magnetometer principles of operation | 191 |
| 10.2.1 | Helium resonance element | 192 |
| 10.2.2 | Helium optical pumping radiation sources | 192 |
| 10.2.3 | Optical pumping of metastable helium | 194 |
| 10.2.4 | Observation of optically pumped helium | 196 |
| 10.2.5 | Observation of magnetic resonance signals in optically pumped helium | 197 |
| 10.3 | Conclusions | 202 |
| 11 | Surface coatings for atomic magnetometry | 205 |
| | <i>S. J. Seltzer, M.-A. Bouchiat, and M. V. Balabas</i> | |
| 11.1 | Introduction and history | 205 |

| | | |
|----------------|--|-----|
| 11.2 | Wall relaxation mechanisms | 208 |
| 11.2.1 | Origin and time dependence of the disorienting interaction | 208 |
| 11.2.2 | Methods of investigation | 209 |
| 11.2.3 | Quantitative interpretation | 212 |
| 11.3 | Coating preparation | 213 |
| 11.4 | Light-induced atomic desorption (LIAD) | 217 |
| 11.5 | Recent characterization methods | 219 |
| 12 | Magnetic shielding | 225 |
| | <i>V. V. Yashchuk, S.-K. Lee, and E. Paperno</i> | |
| 12.1 | Introduction | 225 |
| 12.2 | Ferromagnetic shielding | 225 |
| 12.2.1 | Simplified estimation of ferromagnetic shielding efficiency for a static magnetic field | 226 |
| 12.2.2 | Multilayer ferromagnetic shielding | 227 |
| 12.2.3 | Optimization of permeability: annealing, degaussing, shaking, tapping | 232 |
| 12.2.4 | Magnetic-field noise in ferromagnetic shielding | 235 |
| 12.2.5 | Examples of ferromagnetic shielding systems | 236 |
| 12.3 | Ferrite shields | 238 |
| 12.3.1 | Permeability | 238 |
| 12.3.2 | Fabrication and the effect of an air gap | 239 |
| 12.3.3 | Thermal noise | 240 |
| 12.4 | Superconducting shields | 241 |
| 12.4.1 | Principles | 242 |
| 12.4.2 | Materials and fabrication | 243 |
| 12.4.3 | Image field | 244 |
| Part II | Applications | 249 |
| 13 | Remote detection magnetometry | 251 |
| | <i>S. M. Rochester, J. M. Hight, B. Patton, D. Budker, R. Holzlöhner, and D. Bonaccini Calia</i> | |
| 13.1 | Introduction | 251 |
| 13.2 | A remotely interrogated all-optical ^{87}Rb magnetometer | 252 |
| 13.3 | Magnetometry with mesospheric sodium | 256 |
| 14 | Detection of nuclear magnetic resonance with atomic magnetometers | 265 |
| | <i>M. P. Ledbetter, I. Savukov, S. J. Seltzer, and D. Budker</i> | |
| 14.1 | Introduction | 265 |
| 14.2 | The NMR Hamiltonian | 267 |
| 14.3 | Challenges associated with detection of NMR using atomic magnetometers | 268 |
| 14.4 | Remote detection | 269 |

| | | |
|--------|--|-----|
| 14.5 | Solenoid matching of Zeeman resonance frequencies | 272 |
| 14.6 | Flux transformer | 273 |
| 14.7 | Nuclear quadrupole resonance | 274 |
| 14.8 | Zero-field nuclear magnetic resonance | 275 |
| 14.8.1 | Thermally polarized zero-field NMR J spectroscopy | 275 |
| 14.8.2 | Parahydrogen-enhanced zero-field NMR | 278 |
| 14.8.3 | Zeeman effects on J -coupled multiplets | 281 |
| 14.9 | Conclusions | 282 |
| 15 | Space magnetometry | 285 |
| | <i>B. Patton, A. W. Brown, R. E. Slocum, and E. J. Smith</i> | |
| 15.1 | Introduction | 285 |
| 15.1.1 | Achievements of space magnetometry | 285 |
| 15.1.2 | Challenges unique to space magnetometers | 286 |
| 15.1.3 | Magnetic sensors used in space missions | 287 |
| 15.2 | Alkali-vapor magnetometers in space applications | 287 |
| 15.2.1 | Initial development of Earth's-field alkali magnetometers | 287 |
| 15.2.2 | Sensor design | 288 |
| 15.2.3 | NASA missions employing alkali-vapor magnetometers | 289 |
| 15.3 | Helium magnetometers in space applications | 293 |
| 15.3.1 | Introduction | 293 |
| 15.3.2 | Future helium space magnetometers | 298 |
| 16 | Detection of biomagnetic fields | 303 |
| | <i>A. Ben-Amar Baranga, T. G. Walker, and R. T. Wakai</i> | |
| 16.1 | Sources of biomagnetism | 303 |
| 16.2 | Development of biomagnetic field detection | 304 |
| 16.3 | Medical applications | 308 |
| 16.4 | Magnetocardiography with atomic magnetometers | 310 |
| 16.5 | Magnetoencephalography with an atomic magnetometer | 313 |
| 16.6 | Summary | 316 |
| 17 | Geophysical applications | 319 |
| | <i>M. D. Prouty, R. Johnson, I. Hrvoic, and A. K. Vershovskiy</i> | |
| 17.1 | Airborne magnetometers and gradiometers | 319 |
| 17.2 | Ground magnetometers/gradiometers | 321 |
| 17.3 | Marine magnetometers/gradiometers | 323 |
| 17.4 | Vector magnetometry with optically pumped magnetometers | 324 |
| 17.5 | Earthquake studies | 329 |
| 17.6 | Applications of magnetometers to detecting unexploded ordnance (UXO) | 331 |
| 17.6.1 | Introduction to the problem | 331 |
| 17.6.2 | Using magnetometers for UXO detection | 332 |
| 17.6.3 | Mathematics of UXO detection | 333 |

| | |
|--|-----|
| Part III Broader impact | 337 |
| 18 Tests of fundamental physics with optical magnetometers | 339 |
| <i>D. I. Jackson Kimball, S. K. Lamoreaux, and J. E. Chupp</i> | |
| 18.1 Overview and introduction | 340 |
| 18.2 Searches for permanent electric dipole moments | 341 |
| 18.2.1 Basic experimental setup for an EDM experiment | 344 |
| 18.2.2 Sensitivity to EDMs | 345 |
| 18.2.3 Electric fields and coherence times for various systems | 346 |
| 18.2.4 Magnetometry and comagnetometry in EDM experiments | 349 |
| 18.3 Anomalous spin-dependent forces | 352 |
| 18.3.1 Background | 352 |
| 18.3.2 Experiments | 355 |
| 18.4 <i>CPT</i> and local Lorentz invariance tests | 361 |
| 18.5 Conclusion | 364 |
| 19 Nuclear magnetic resonance gyroscopes | 369 |
| <i>E. A. Donley and J. Kitching</i> | |
| 19.1 Introduction | 369 |
| 19.2 NMR frequency shifts and relaxation | 373 |
| 19.2.1 Spin exchange | 374 |
| 19.2.2 Quadrupole surface frequency shifts | 375 |
| 19.2.3 General wall relaxation | 377 |
| 19.2.4 Magnetic-field gradients | 377 |
| 19.2.5 Noble-gas self-relaxation | 378 |
| 19.3 Alkali shifts and relaxation mechanisms | 379 |
| 19.4 Two-spin NMR gyroscope | 379 |
| 19.5 Comagnetometer | 381 |
| 19.6 Miniaturization | 383 |
| 19.7 Conclusion | 383 |
| 20 Commercial magnetometers and their application | 387 |
| <i>D. C. Hovde, M. D. Prouty, I. Hrvoic, and R. E. Slocum</i> | |
| 20.1 Introduction | 387 |
| 20.2 Specifications | 388 |
| 20.2.1 Noise | 388 |
| 20.2.2 Resolution | 391 |
| 20.2.3 Sensitivity | 391 |
| 20.2.4 Sample rate and cycle time | 392 |
| 20.2.5 Bandwidth | 392 |
| 20.2.6 Absolute error and drift | 393 |
| 20.2.7 Gradient tolerance | 394 |
| 20.2.8 Dead zones | 395 |
| 20.2.9 Heading error | 395 |
| 20.2.10 Range of measurement | 397 |

| | | |
|--------|---|-----|
| 20.2 | History of commercial magnetometry | 398 |
| 20.3.1 | Fluxgate magnetometers | 398 |
| 20.3.2 | SQUID magnetometers | 399 |
| 20.3.3 | Proton-precession and Overhauser magnetometers | 399 |
| 20.3.4 | Alkali metal magnetometers: rubidium, cesium, and potassium | 401 |
| 20.3.5 | Helium-3 and helium-4 magnetometers | 402 |
| 20.4 | Military applications | 403 |
| 20.5 | Anticipated improvements | 404 |
| | <i>Index</i> | 406 |