


# Lecture Notebook



*for* Campbell and Farrell's  
**Biochemistry**  
Fourth Edition

Mary K. Campbell  
Shawn O. Farrell

## Table of Contents

| <b>Image Number</b> | <b>Figure</b> | <b>Description</b>   |
|---------------------|---------------|--|
| 1–3                 | 1.10          | A comparison of (a) an animal cell, (b) a plant cell, and (c) a prokaryotic cell |
| 4                   | Table 1.1     | Functional groups of biochemical importance                                      |
| 5                   | 2.9a          | Titration of acetic acid with NaOH   |
| 6                   | 2.9b          | Relative abundance of acetic acid and acetate ion during a titration             |
| 7–8                 | 2.11          | Titration curve and buffering action   |
| 9                   | Table 2.8     | Acid and base forms of some buffers  |
| 10                  | 3.1           | Amino acid structure   |
| 11–14               | 3.4           | Amino acids and their structures   |
| 15–16               | 3.7           | Titration curves of amino acids  |
| 17                  | Table 3.1     | Common amino acids and their abbreviations                                       |
| 18                  | Table 3.2     | PKa values of common amino acids   |
| 19                  | 4.1           | Angles of the polypeptide chain  |
| 20                  | 4.2           | The $\alpha$ -helix  |
| 21                  | 4.3           | $\beta$ -pleated sheet   |
| 22                  | 4.12          | Forces that stabilize tertiary structure in proteins                             |
| 23                  | 4.18          | Denaturation of proteins   |
| 24–25               | A.5           | Gel filtration chromatography  |
| 26                  | A.6           | Affinity chromatography  |
| 27                  | A.8           | Ion exchange chromatography using a cation exchanger                             |
| 28                  | A.13          | Strategy for determining the primary structure of a given protein                |
| 29                  | A.18          | Sequencing of peptides by the Edman method                                       |
| 30                  | 5.1           | Activation energy profiles   |
| 31                  | 5.10          | Lineweaver-Burk double-reciprocal plot   |
| 32                  | 5.12          | Lineweaver-Burk double-reciprocal plot for competitive inhibition                |
| 33                  | 5.13          | Lineweaver-Burk double-reciprocal plot for noncompetitive inhibition             |
| 34                  | 6.4           | Monod-Wyman-Changeux model   |
| 35                  | 6.5           | Monod-Wyman-Changeux or concerted model  |
| 36                  | 6.7           | Sequential model of cooperative binding of substrate to allosteric enzyme        |
| 37                  | 6.8           | Phosphorylation of the sodium/potassium pump                                     |
| 38                  | 6.9           | Glycogen phosphorylation activity  |
| 39–40               | 6.13          | The mechanism of chymotrypsin action   |
| 41                  | 7.5           | Structures of some phosphoacylglycerols  |
| 42                  | 7.8           | Ganglioside structures   |
| 43                  | 7.11          | Lipid bilayer asymmetry  |
| 44                  | 7.18          | Fluid mosaic model for membrane structure  |
| 45                  | 7.24          | The sodium-potassium pump  |
| 46                  | 7.26          | Secondary active transport   |
| 47–48               | 8.4           | Commonly occurring nucleotides   |
| 49                  | 8.5           | A fragment of an RNA chain   |
| 50                  | 8.6           | A portion of a DNA chain   |

| <b>Image Number</b> | <b>Figure</b> | <b>Description</b>  |
|---------------------|---------------|---|
| 51                  | 8.7           | The double helix  |
| 52                  | 8.11          | A Z-DNA section in the middle of a B-DNA  |
| 53                  | 8.15          | A model for the action of bacterial DNA gyrase  |
| 54                  | 8.19          | Types of RNA  |
| 55                  | 9.2           | Semiconservative replication  |
| 56                  | 9.4           | Bi-directional replication of DNA in prokaryotes and eukaryotes                               |
| 57                  | 9.5           | The semidiscontinuous model for DNA replication   |
| 58                  | 9.10          | General features of the replication fork  |
| 59                  | 9.12          | The 5' → 3' exonuclease activity of DNA polymerase I  |
| 60                  | 9.16          | Base excision repair  |
| 61                  | 9.19          | Model for the initiation of the DNA replication cycle in eukaryotes                           |
| 62                  | 10.3          | The basic order of events in prokaryotic transcription initiation and elongation              |
| 63                  | 10.4          | Topoisomerases remove supercoils that would form ahead of and behind the transcription bubble |
| 64                  | 10.6          | Transcription termination by the rho factor   |
| 65–66               | 10.9          | The lacI gene produces a protein that represses the lac operon                                |
| 67                  | 10.12         | Basic control mechanisms seen in the expression of genes                                      |
| 68                  | 10.13         | The trp operon of E. coli.  |
| 69                  | 10.18         | The order of events of transcription  |
| 70                  | 10.27         | Amino acid sequence comparisons of several DNA binding proteins                               |
| 71                  | 10.30         | Posttranscriptional modification of tRNA precursor  |
| 72                  | 10.34         | Splicing of mRNA precursors   |
| 73                  | 10.35         | Organization of the fast skeletal muscle troponin T gene                                      |
| 74–75               | 11.3          | The filter-binding assay for elucidation of the genetic code                                  |
| 76                  | 11.5          | Various base-pairing alternatives   |
| 77                  | 11.9          | Formation of the N-formylmethionine-tRNA  |
| 78                  | 11.12         | A summary of the steps in chain elongation  |
| 79                  | 11.15         | The events in peptide chain termination   |
| 80                  | 11.17         | Simultaneous protein synthesis on polysomes   |
| 81                  | 11.20         | The three stages in the initiation of translation in eukaryotic cells                         |
| 82                  | 11.25         | The outcome of infection of cells by simian virus 40 depends on the nature of the cells       |
| 83                  | Table 11.1    | The genetic code  |
| 84                  | B.4           | Hydrolysis of DNA by restriction endonucleases  |
| 85                  | B.5           | Restriction endonuclease EcoRI cleaves double-stranded DNA                                    |
| 86                  | B.6           | The cloning of a virus  |
| 87                  | B.8           | The cloning of human DNA fragments with a viral vector  |
| 88                  | B.9           | Selecting for recombinant DNA in a bacterial plasmid  |
| 89                  | B.12          | The pUC series of plasmids  |
| 90                  | B.13          | Clone selection via blue/white screening  |
| 91                  | B.15          | Synthesis of insulin in humans  |
| 92                  | B.18          | Gene therapy in bone marrow cells   |

| <b>Image Number</b> | <b>Figure</b> | <b>Description</b>   |
|---------------------|---------------|--|
| 93                  | B.19          | Human gene therapy via retroviruses  |
| 94                  | B.23          | Steps involved in the construction of a DNA library  |
| 95                  | B.25          | Selecting a desired clone from a DNA library   |
| 96                  | B.26          | The polymerase chain reaction  |
| 97                  | B.27          | The Southern blot  |
| 98                  | B.29          | Restriction fragment length polymorphisms  |
| 99                  | B.34          | Using a reporter gene  |
| 100                 | 12.4          | Catabolism and anabolism   |
| 101                 | 12.6          | NADH and NAD <sup>+</sup>  |
| 102                 | 12.14         | Anabolism and catabolism   |
| 103                 | 13.2          | An-aldose and a ketose   |
| 104                 | 13.4          | Stereochemical relationships among monosaccharides   |
| 105                 | 13.8          | A comparison of the Fischer, complete Haworth, and abbreviated Haworth structures of $\alpha$ - and $\beta$ -D-glucose |
| 106                 | 14.1          | One molecule of pyruvate is converted to two molecules of pyruvate   |
| 107                 | 14.2          | The pathway of glycolysis  |
| 108                 | 14.3          | The first phase of glycolysis  |
| 109                 | 14.7          | The second phase of glycolysis   |
| 110                 | Table 14.1    | The reactions of glycolysis and their standard free energy changes   |
| 111                 | 15.5          | Glycogen phosphorylase activity is subject to allosteric control and covalent modification                             |
| 112                 | 15.6          | The pathways of gluconeogenesis and glycolysis   |
| 113                 | 15.12         | The Cori cycle   |
| 114                 | 15.15         | The oxidative reactions of the pentose phosphate pathway   |
| 115                 | 15.16         | The nonoxidative reactions of the pentose phosphate pathway  |
| 116                 | 15.17         | Relationship between the pentose phosphate pathway and glycolysis  |
| 117                 | 16.1          | The citric acid cycle and catabolism   |
| 118                 | 16.3          | An overview of the citric acid cycle   |
| 119                 | 16.4          | The mechanism of the pyruvate dehydrogenase reaction   |
| 120                 | 16.7          | Control points in the conversion of pyruvate to acetyl-CoA and in the citric acid cycle                                |
| 121                 | 16.8          | The glyoxylate cycle   |
| 122                 | 16.9          | A summary of catabolism  |
| 123                 | 16.11         | Transfer of the starting materials of glyconeogenesis from the mitochondrion to the cytosol                            |
| 124                 | 16.12         | Transfer of the starting materials of lipid anabolism from the mitochondrion to the cytosol                            |
| 125                 | 16.14         | A summary of anabolism   |
| 126                 | Table 16.2    | The energetics of conversion of pyruvate to carbon dioxide   |
| 127                 | 17.7          | Respiratory complexes in the inner mitochondrial membrane  |
| 128                 | 17.11         | A model for the components of ATP synthase   |
| 129                 | 17.13         | The creation of a proton gradient in chemiosmotic coupling   |

| <b>Image Number</b> | <b>Figure</b> | <b>Description</b>   |
|---------------------|---------------|--|
| 130                 | 17.21         | The glycerol phosphate shuttle   |
| 131                 | 17.22         | The malate-aspartate shuttle   |
| 132                 | Table 17.1    | Standard reduction potentials for biological half-reactions  |
| 133                 | Table 17.2    | Energetics of electron transport reactions   |
| 134                 | Table 17.3    | Yield of ATP from glucose oxidation  |
| 135                 | 18.3          | Liberation of fatty acids from triacylglycerols  |
| 136                 | 18.5          | The role of carnitine in the transfer of acyl groups to the mitochondrial matrix                   |
| 137                 | 18.6          | The $\beta$ -oxidation of saturated fatty acids  |
| 138                 | 18.8          | The oxidation of fatty acid containing an odd number of carbon atoms                               |
| 139                 | 18.10         | The oxidation pathway for polyunsaturated fatty acids  |
| 140–141             | 18.15         | The first cycle of palmitate synthesis   |
| 142–143             | 18.19         | Pathways for the biosynthesis of triacylglycerols  |
| 144                 | Table 18.1    | The balance sheet for oxidation of one molecule of stearic acid                                    |
| 145                 | 19.5          | Electron flow in Photosystems I and II   |
| 146                 | 19.8          | Cyclic electron flow coupled to photophosphorylation in Photosystem I                              |
| 147–148             | 19.9          | Molecular events that take place at the photosynthetic reaction center of <i>Rhodospseudomonas</i> |
| 149                 | 19.11         | The relationship between photophosphorylation and the proton gradient                              |
| 150                 | 19.12         | The components of the electron transport chain of the thylakoid membrane                           |
| 151                 | 19.13         | The two possible electron transfer pathways in a photosynthetic anaerobe                           |
| 152                 | 19.14         | The main features of the Calvin cycle  |
| 153                 | 19.20         | The complete Calvin cycle  |
| 154                 | 19.21         | The C <sub>4</sub> pathway   |
| 155                 | 20.5          | Families of amino acids based on biosynthetic pathways   |
| 156                 | 20.6          | The relationship between amino acid metabolism and the citric acid cycle                           |
| 157                 | 20.17         | The urea cycle   |
| 158                 | 20.18         | The urea cycle and some of its links to the citric acid cycle                                      |
| 159                 | C.4           | Feedback control in the biosynthesis of amino acids of the aspartate family                        |
| 160                 | C.9           | Histidine biosynthesis   |
| 161                 | C.10          | The biosynthesis of porphyrins   |
| 162                 | 21.1          | Block diagram of intermediary metabolism   |
| 163                 | 21.4          | Hormonal control system  |
| 164                 | 21.5          | Activation of adenylyl cyclase by heterodimeric G proteins   |
| 165                 | 21.12         | A two-stage process leads to the growth and differentiation of T cells                             |